

## A BROADBAND STUDY OF THE EMISSION FROM THE COMPOSITE SUPERNOVA REMNANT MSH 11–62

PATRICK SLANE,<sup>1</sup> JOHN P. HUGHES,<sup>2</sup> TEA TEMIM,<sup>3</sup> ROMAIN ROUSSEAU,<sup>4</sup> DILLON FOIGHT,<sup>1</sup> B. M. GAENSLER,<sup>5</sup>  
STEFAN FUNK,<sup>6</sup> MARIANNE LEMOINE-GOUMARD,<sup>4</sup> JOSEPH D. GELFAND,<sup>7</sup> DAVID A. MOFFETT,<sup>8</sup> RICHARD G. DODSON,<sup>9</sup> AND  
JOSEPH P. BERNSTEIN,<sup>10,11</sup>

*Draft version September 17, 2011*



### ABSTRACT

MSH 11–62 (G291.1–0.9) is a composite supernova remnant for which radio and X-ray observations have identified the remnant shell as well as its central pulsar wind nebula. The observations suggest a relatively young system expanding into a low density region. Here we present a study of MSH 11–62 using observations with the *Chandra*, *XMM*, and *Fermi* observatories, along with radio observations from the Australia Telescope Compact Array (ATCA). We identify a compact X-ray source that appears to be the putative pulsar that powers the nebula, and show that the X-ray spectrum of the nebula bears the signature of synchrotron losses as particles diffuse into the outer nebula. Using data from the *Fermi* LAT, we identify  $\gamma$ -ray emission originating from MSH 11–62. With density constraints from the new X-ray measurements of the remnant, we carry out models for the evolution of the composite system in order to constrain the properties of the underlying pulsar and the origin of the  $\gamma$ -ray emission.

*Subject headings:* radiation mechanisms: non-thermal — supernova remnants: individual: MSH 11–62  
— X-rays: ISM — stars: neutron

### 1. INTRODUCTION

The creation of an energetic pulsar in a supernova explosion often results in the formation of a composite supernova remnant (SNR) that can be characterized by emission from the remnant, an associated pulsar wind nebula (PWN), and the neutron star (NS) itself. Recent studies of such systems have resulted in discoveries of new pulsars, constraints on the spectra of particles injected from the pulsars into their nebulae, and evidence for interactions between the SNR reverse shock (RS) and the PWN. Broadband observations of composite SNRs have been crucial for improving our understanding of the detailed evolution of these systems, particularly with the vast improvements in capabilities provided by current X-ray and  $\gamma$ -ray observatories.

MSH 11–62 (Mills, Slee, & Hill 1961) is a centrally-bright SNR whose properties strongly suggest the presence of a central nebula powered by an active pulsar. Radio observations reveal a remnant with a distinct bar-like central structure whose flat radio spectrum ( $\alpha_r = 0.29 \pm 0.05$ , where  $S_\nu \propto \nu^{-\alpha_r}$ ) is similar to that observed for PWNe (Roger et al. 1986). Polarization measurements at 5 and 8.4 GHz indicate that the magnetic field in the central regions of the SNR is aligned with the long axis of the bar.

The distance to MSH 11–62 is poorly known. Limits on HI absorption indicate a distance greater than 3.5 kpc (Moffett, Gaensler, & Green 2001; Moffett et al. 2002). We adopt a fiducial distance of 5 kpc in derived quantities below, and scale values to  $d_5$ , the distance in units of 5 kpc. Based on the radio synchrotron spectrum, the minimum-energy magnetic field in the central nebula is  $B \sim 65d_5^{-2/7} \mu\text{G}$  (Harrus, Hughes, & Slane 1998).

X-ray emission from MSH 11–62 was first identified by Wilson (1986) using the *Einstein Observatory*. X-ray observations with the *ASCA* observatory reveal a spectrum best described by a combination of thermal and non-thermal models (Harrus, Hughes, & Slane 1998). The hard X-ray emission is spatially concentrated in the central regions, indicating the presence of a PWN within the remnant, although an 11 hr observation at 1.4 GHz failed to detect any pulsations from this region, down to a sensitivity limit of 0.028 mJy assuming a 10% duty cycle<sup>9</sup>. The derived spectral index for the X-ray power law component is  $\Gamma = 2.0_{-0.3}^{+0.1}$  and the associated luminosity is  $L_x(0.2 - 4 \text{ keV}) \sim 6 \times 10^{33} d_5^2 \text{ erg s}^{-1}$ . The best-fit temperature of the thermal component, presumably associated with emission from shock-heated gas in the SNR shell, is  $kT \sim 0.8 \text{ keV}$ . These observations clearly imply that MSH 11–62 harbors a young pulsar that powers a central PWN. *Chandra* observations, which we discuss in detail below, reveal a compact X-ray source that is almost certainly the pulsar counterpart (Harrus et al. 2002; Kargaltsev & Pavlov 2008).

Observations with the *Compton Gamma-Ray Observatory* reveal the source 3EG J1102–6103, for which MSH 11–62 falls within the position error circle. The flux is  $F(> 100 \text{ MeV}) = (3.3 \pm 0.6) \times 10^{-7} \text{ photons cm}^{-2} \text{ s}^{-1}$ , with a power law index  $\Gamma = 2.5 \pm 0.2$  (Hartman et al. 1999). This source was suggested as a  $\gamma$ -ray counterpart to MSH 11–62 (Sturmer & Dermer 1995), but was also subsequently suggested as the counterpart to the nearby young pulsar J1105–6107 (Kaspi et al. 1997). Given the

<sup>1</sup> Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138-1516; slane@cfa.harvard.edu.

<sup>2</sup> Dept. of Physics and Astronomy, Rutgers Univ., Piscataway, NJ 08854-8019; jph@physics.rutgers.edu.

<sup>3</sup> NASA Goddard Space Flight Center - Code 662, Greenbelt, MD 20771

<sup>5</sup> Sydney Institute for Astronomy, School of Physics A29, The University of Sydney, NSW 2006, Australia; bryan.gaensler@sydney.edu.au

<sup>8</sup> Dept. of Physics, Furman Univ., Greenville, SC 29613; david.moffett@furman.edu.

<sup>9</sup> International Centre for Radio Astronomy Research University of Western Australia; richard.dodson@icrar.org.

<sup>10</sup> High Energy Physics Division, Argonne National Laboratory, Argonne, IL 60439; jpbernst@anl.gov.

<sup>11</sup> Department of Astronomy, University of Michigan, Ann Arbor, MI 48109

<sup>9</sup> F. Camilo, private communication.




FIG. 1.— ATCA 1.4 GHz image of MSH 11–62. The remnant is characterized by a faint shell with a bright central bar. Contours levels are 3.6 and 1.5 mJy beam<sup>-1</sup>. The cross marks the position of the X-ray point source described in Section 3.

large uncertainty in the position of the  $\gamma$ -ray source, the origin of its emission remained uncertain.

Here we report on observations with the ATCA, the *Chandra X-ray Observatory*, the *XMM-Newton X-ray Observatory*, and the *Fermi Gamma-Ray Space Telescope* that provide sufficient sensitivity and resolution to identify the neutron star in MSH 11–62, and to provide a much-improved characterization of the properties of this composite system. In Sections 2 we discuss our ATCA observations. In Section 3 we summarize the *Chandra* and *XMM* observations, with detailed investigations of the PWN, neutron star, and SNR shell. In Section 4 we discuss observations with the *Fermi LAT*, and in Section 5 we present discussions of the results along with our conclusions.

## 2. RADIO OBSERVATIONS

### 2.1. Observations and Data Reduction

Radio observations of MSH 11–62 were carried out with the ATCA, a 6-element interferometer located near Narrabri, New South Wales, Australia.

Observations were carried out on three separate dates using three different array configurations: on 1999 Feb 10 (750C array), 1999 Feb 22 (6C array), and 1999 Mar 24 (1.5B array). In all cases, radio continuum data were recorded in all four polarizations, over a bandwidth of 128 MHz centered at 1384 MHz. At each epoch, MSH 11–62 was observed for 8–11 hours spread over a 12-hour synthesis, combined with regular short observations of the extragalactic source PKS B1036–697 to calibrate atmospheric gain variations and polarization leakages. The bright source PKS B1934–638 was also observed at each epoch to provide an overall flux calibration.

Following editing of the data and calibration in the MIRIAD package using standard techniques, the visibility data-set was imaged using multi-frequency synthesis and uniform weighting, and then deconvolved with a maxi-




FIG. 2.— *Chandra* image of MSH 11–62. Colors correspond to different X-ray energy bands: red = 0.75 - 1.45 keV; green = 1.45 - 2.58 keV; blue = 2.58 - 6.0 keV. The SNR shell is evident as soft (red) X-ray emission (the portion of the shell in the northeast falls outside the detector field of view). The central bar and associated compact source are white in color, corresponding to harder emission.

mum entropy algorithm. The final total intensity image has an angular resolution of  $8''.8 \times 6''.6$ , with the major axis oriented at a position angle  $3^\circ$  east of north, and an RMS sensitivity of 0.1–0.2 mJy beam<sup>-1</sup>. Preliminary investigations of these observations have been presented by Moffett, Gaensler, & Green (2001) and Moffett et al. (2002).

### 2.2. Radio Structure and Properties

The ATCA 1.4 GHz image of MSH 11–62 (Figure 1) reveals a roughly circular morphology with a bright central bar-like structure that corresponds with the flat-spectrum region identified by Roger et al. (1986). Partial limb-brightening is observed along the northwestern and southeastern boundaries of the SNR shell. Considerable filamentary structures are observed both along the limb and in the SNR interior. The bright central bar reveals several regions of enhanced radio emission, with the brightest feature located near the southeastern end of the structure (at an approximate position of  $11^{\text{h}}11^{\text{m}}48^{\text{s}}, -60^{\circ}39'27''$ ). The structure of the central bar is similar to that observed in several other PWNs, including G11.2–0.3 (Tam et al. 2002), N157B (Lazendic et al. 2000), and G328.4+0.2 (Johnston, McClure-Griffiths, & Koribalski 2004; Gelfand et al. 2007). Its apparent similarity to the PWN in G328.4+0.2, in particular, may suggest that the nebula has been compressed by the SNR reverse shock (RS), which may have propagated more quickly from the NW/SE directions, where the SNR limbs show slight brightness enhancements.

The total 1.4 GHz flux from the SNR is 14 Jy, with 4.8 Jy associated with the central nebula.

## 3. X-RAY OBSERVATIONS

A 50 ks *Chandra* ACIS observation of MSH 11–62 was performed on 2002 April 8 (ObsID 2782). The central region of the SNR was placed on the S3 chip of the de-




FIG. 3.— *Chandra* image of PWN in MSH 11–62, along with regions used for extracting spectral index along the nebula (see Table 1). The image has been smoothed with a Gaussian of 5 arcsec width. The pulsar candidate is the bright central source between regions 1 and 4.

detector, and portions of the shell were imaged on the adjacent chip S2. The extent of the SNR (with a radius of roughly 6 arcmin) is considerably larger than the field covered by the S3 chip, which resulted in the northeastern regions of the remnant falling outside of the detector field of view.

MSH 11–62 was observed for 40 ks with *XMM*, on 2002 February 6 (ObsID 0051550101). The MOS2 and pn detectors were operated in full frame mode with the MEDIUM filters, providing full coverage of the SNR, while MOS1 was operated in timing mode with the THIN filter.

### 3.1. Observations and Data Reduction

The *Chandra* data were reduced using standard screening prescriptions. All data were reduced using tools from *CIAO* 3.2.2. After screening, the remaining exposure time was 49.4 ks.

The *Chandra* observations show hard emission extended along the radio bar (Figure 2), although the emission does not extend as far to the NE as the radio bar. There is also a clear shell of soft X-ray emission that corresponds well with the morphology of the radio SNR, including a small intensity enhancement along the position of the western radio limb-brightening, and a large underluminous region to the northwest of the radio bar.

Of particular significance in the *Chandra* image is the presence of numerous point sources of X-ray emission along the line of sight to MSH 11–62. These appear to be associated with the open cluster Tr 18, located at a distance of  $\sim 1.5$  kpc (Vazquez & Feinstein 1990).

We have analyzed spectra from both the central X-ray bar and the SNR shell using background regions selected from regions just outside the boundaries of the remnant. Weighted response matrix (RMF) and ancillary response




FIG. 4.— *Chandra* ACIS X-ray spectra for PWN (black) and NS (red) with absorbed power law fits.

(ARF) files were created for the extended regions, and spectral modeling was then carried out using XSPEC version 11.3.2ag. Point sources within the regions of interest were excluded from the spectral extraction.

The *XMM* data were reduced using SAS Version 10.0. Standard screening resulted in cleaned exposures of 35.6 (29.2) ks for MOS2 (pn).

### 3.2. Central Bar

The central X-ray bar is aligned with the radio bar, and extends to the edge of the radio in the southwest, but the brightness falls off more quickly than the radio emission in the northeast direction. There is a bright X-ray source located at the position of the brightest radio emission, likely representing the pulsar that powers the nebula. The source, located at  $11^{\text{h}}11^{\text{m}}48^{\text{s}}.62, -60^{\circ}39'26''.2$  (and hereafter identified as CXOU J111148.6-603926), is roughly centered on the X-ray nebula. A fainter point source is detected at the southwestern edge of the radio bar, with coordinates  $11^{\text{h}}11^{\text{m}}44^{\text{s}}.5, -60^{\circ}40'02''.8$ . The source is positionally coincident with the  $V = 10.8$  star Tycho 8959 255 1, and is unrelated to the nebula.

The *Chandra* spectrum of the X-ray bar is well-described by a power law model with  $N_H \approx 7 \times 10^{21} \text{ cm}^{-2}$  and  $\Gamma \approx 1.7$  (Table 1, Figure 3), securing the PWN interpretation for this structure. Here we have used an elliptical background region on S3, adjacent to the PWN but within the SNR, to account for both sky and instrumental background as well as thermal emission from the SNR. An independent fit using the *XMM* pn data yields virtually identical results. The luminosity (0.5 – 10 keV) is  $1.1 \times 10^{34} d_5^2 \text{ erg s}^{-1}$ , about a factor of two higher than the value determined from *ASCA* observations (Harrus, Hughes, & Slane 1998), but far lower than the value of  $\sim 8 \times 10^{34} d_5^2 \text{ erg s}^{-1}$  cited by Kargaltsev & Pavlov (2008).

To search for spectral variations in the PWN, we ex-

TABLE 1  
SPECTRAL FIT PARAMETERS

Region	Data	Parameter	Value
PWN (whole)	CXO	$N_H$	$(6.7 \pm 0.7) \times 10^{21} \text{ cm}^{-2}$
		$\Gamma$	$1.8 \pm 0.1$
		$F_x^a$	$3.5 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$
PWN Region	CXO	$N_H$	(fixed)
		$\Gamma$	$1.7 \pm 0.1$
			$1.8 \pm 0.1$
			$2.2 \pm 0.1$
			$1.5 \pm 0.1$
			$1.7 \pm 0.1$
NS	CXO	$N_H$	(fixed)
		$\Gamma$	$1.2 \pm 0.2$
		$F_x^a$	$3.7 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$
SNR (whole)	XMM	$N_H$	(fixed)
		$kT_1$	$1.3^{+0.5}_{-0.2} \text{ keV}$
		$n_e t$	$(1.5 \pm 0.2) \times 10^{10} \text{ s cm}^{-3}$
		$kT_2$	$2.8 \pm 0.4 \text{ keV}$
		$F_1^a$	$2.1 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$
		$F_2^a$	$3.0 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$

NOTE. — a) Unabsorbed 0.5 - 10.0 keV flux.

tracted spectra from rectangular regions along the nebula, as shown in Figure 3. We fixed the column density to that obtained for the entire PWN, and then fit each spectrum to a power law model. The results, summarized in Table 1, show a spectral steepening with distance from the central neutron star. Such steepeng is also observed in G21.5–0.9 (Slane et al. 2000), 3C 58 (Slane et al. 2004) and other PWNe, and is understood to be associated with synchrotron aging of the particles as they diffuse from the injection point to the outer nebula (Kennel & Coroniti 1984), although models matching the exact radial behavior of the spectral index remain elusive (e.g. Reynolds 2003).

### 3.3. Compact Source

Our *Chandra* observations obtain 820 background-subtracted counts from within a circle with a radius of 3.3 arcsec centered on the compact source CXOU J111148.6-603926. The associated count rate of  $1.6 \times 10^{-2} \text{ cnt s}^{-1}$  yields negligible pileup in the image. The radial profile of the compact source image is well-described by the point spread function of the telescope, with no evidence of a surrounding jet/torus structure as is seen in many other such systems, although evidence for filamentary structure is seen on larger scales (see Figure 3).

The spectrum of the compact source is well-described by a power law with the column density fixed at the value obtained from fits of the entire PWN, and with  $\Gamma = 1.2 \pm 0.2$ . The unabsorbed flux is  $F_x(0.5-10 \text{ keV}) = 3.7 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$  indicating a luminosity  $L_x(0.5-10 \text{ keV}) = 1.1 \times 10^{33} d_5^2 \text{ erg s}^{-1}$ , similar to that measured for a number of other young pulsars with bright PWNe (see Kargaltsev & Pavlov 2008).




FIG. 5.— *XMM* MOS2 image of MSH 11-62, with radio contours from Figure 1. The image has been exposure-corrected, and red/green/blue corresponds to the energy band 0.6-1.5/1.5-2.5/2.5-5.0 keV. The large circle indicates the region from which the SNR spectrum was extracted. Events from the inner ellipse were ignored to eliminate the contribution from the PWN, and those from the small circles were ignored to reduce point source contributions. The same inner ellipse was used to extract the spectrum of the PWN.

### 3.4. SNR Shell

To investigate the X-ray properties of the SNR shell, we extracted spectra of the entire SNR from both the *XMM* pn and MOS2 detectors (see Figure 6). Emission from the PWN was eliminated from the spectra, as was emission from point sources in the field. Weighted response files were generated for each spectrum, and background spectra were obtained from regions outside the SNR shell in each detector.

The spectra were fit simultaneously to an absorbed nonequilibrium ionization planar shock model (VPSHOCK in XSPEC), with the column density fixed at the best-fit value for the PWN. We find a temperature  $kT = 1.5^{+0.5}_{-0.2} \text{ keV}$  and an ionization timescale  $n_e t = (2.9 \pm 0.9) \times 10^{10} \text{ s cm}^{-3}$ , with evidence for mild overabundances of Ne, Mg, and Si. The fit requires a second, harder component, which can be fit with a power law, but it is best fit with second thermal component (NEI model in XSPEC) with  $kT = 2.8 \pm 0.4 \text{ keV}$  and cosmic abundances.

Based on the volume emission measure for the lower-temperature component, and assuming a thin-shell morphology for the SNR, with a shell thickness of  $R/12$  corresponding to a shock compression ratio of 4, the estimated preshock density is  $n_H = 0.04 \eta^{1/2} d_5^{-1/2} f^{-1/2} \text{ cm}^{-3}$ , where  $\eta$  is the fraction of the X-ray emission that is not associated with point sources in the field, and  $f$  is the fraction of the remnant shell actually seen in X-rays. We note that the higher temperature component provides very nearly the same density estimate.

The mass swept up by the SNR under such a scenario is then  $M_{sw} = 0.5 \eta^{1/2} f^{-1/2} d_5^{5/2}$ . Given that the system morphology appears to show evidence that the PWN has undergone an interaction with the SNR reverse shock, this low value for the swept-up mass would appear to suggest a rather low ejecta mass for the supernova, as we




FIG. 6.— *XMM* spectra from the SNR shell of MSH 11–62, with pn data shown in black and MOS2 data shown in red. The best-fit VPSHOCK model is shown as a histogram.

discuss below.

#### 4. GAMMA-RAY OBSERVATIONS

##### 4.1. *Fermi* LAT

We investigated  $\gamma$ -ray events acquired from the region surrounding MSH 11–62 with the *Fermi*-LAT during the period 2008 August 5 to 2011 March 14. Standard event selection was applied, using “Source” class events with zenith angles less than 100 degrees to minimize the portion of the Earth limb in the LAT field of view (Abdo et al. 2009). The “Pass 7 version 6” instrument response functions were used, along with standard analysis tools available from the *Fermi* Science Support Center (version v9r21p0).

Standard background models were used to account for both Galactic and extragalactic diffuse emission as well as instrumental background. Contributions from neighboring sources identified in the two-year *Fermi*-LAT Second Source Catalog (Abdo et al. 2011) were included in the analysis. The mapcube file *ring\_2year\_P76\_v0.fits* was used to describe the Galactic  $\gamma$ -ray emission, and the isotropic component is modeled using the *isotrop\_2year\_P76\_source\_v0.txt* table. Data analysis details follow those in Castro & Slane (2010) and Grondin et al. (2011).

Using a binned maximum likelihood analysis (the *gt*-like routine) yielded a Test Statistic (TS) corresponding to a significance of  $\sim 21\sigma$  (Mattox et al. 1996). The corresponding TS map is shown in Figure 7, along with (blue) contours from the ATCA observation of MSH 11–62. White contours indicate significance levels of  $\sim 13, 15,$  and  $17\sigma$  for the LAT data. We also indicate the positions of the three known pulsars within a  $45'$  radius from the centroid of the LAT source. If each of these pulsars converts its entire spin-down power into  $\gamma$ -rays in the LAT band, they could account for no more than




FIG. 7.— TS map of *Fermi* LAT emission from MSH 11–62. Blue contours correspond to the radio emission, and white contours represent the  $13, 15,$  and  $17\sigma$  significance levels of the flux. Positions of nearby pulsars with  $\dot{E} > 10^{32}$  erg  $s^{-1}$  are indicated (see discussion in text).

6% of the observed flux, and we thus ignore their contributions in the subsequent analysis. The nearest pulsar with a high  $\dot{E}/d^2$  ratio is PSR J1105–6107, located  $55'$  from the centroid of the  $\gamma$ -ray emission, well beyond the border of Figure 7; we consider an association with this source to be very unlikely.

The source spectrum was extracted using both front and back events, over the energy range 0.1–100 GeV. The spectrum is shown in Figure 8, and is well-described by a power law with  $\Gamma = 1.2 \pm 0.2$  accompanied by an exponential cutoff with  $E_{cut} = 3.2 \pm 0.9$  GeV. The associated flux is  $F(\geq 100 \text{ MeV}) = (6.3 \pm 1.2 \pm 3.2) \times 10^{-8}$  photons  $cm^{-2} s^{-1}$  [or  $(8.0 \pm 0.6 \pm 4.2) \times 10^{-11}$  erg  $cm^{-2} s^{-1}$ ], where the quoted uncertainties are statistical and systematic, respectively.

Two main systematic uncertainties can affect the LAT flux estimation for a point source: uncertainties on the Galactic diffuse background and on the effective area. The dominant uncertainty at low energy comes from the Galactic diffuse emission, which we estimated by changing the normalization of the Galactic diffuse model artificially by 6% as done in Abdo et al. (2010). The second systematic is estimated by using modified IRFs whose effective areas bracket those of our nominal Instrument Response Function (IRF). These “biased” IRFs are defined by envelopes above and below the nominal energy dependence of the effective area by linearly connecting differences of (10%, 5%, 20%) at  $\log(E)$  of (2, 2.75, 4), respectively.

The source 2FGL J1112.1–6040, from the LAT 2<sup>nd</sup> year catalog, was identified with a possible association with MSH 11–62 (Abdo et al. 2011). The 2FGL flux agrees with our measurements as well, but both are significantly lower than that of 3EG J1102–6103, whose quoted spectral index is also significantly steeper than that for MSH 11–62. These differences are presumably associated with the previously-unrecognized spectral cutoff.

We have searched for possible extension of the source




FIG. 8.— *Fermi* LAT spectrum of MSH 11–62. Black error bars correspond to statistical errors, while red error bars include systematic errors as well. The solid curve corresponds to the best-fit model for a power law with an exponential cutoff. The dashed magenta curve corresponds to a pion-decay model assuming a power law proton spectrum with an exponential cutoff. See text for discussion.




FIG. 9.— Upper: Time evolution of the PWN magnetic field. Lower: Evolution of the SNR (blue) and PWN (red) radii as a function of time. The observed radius of MSH 11–62 is indicated, and the vertical line indicates the age at which this radius is reached. See text for model details.

using the routine POINTLIKE, comparing the  $TS$  of the extended hypothesis to the  $TS$  for a point source hypothesis  $TS_p$ , defining  $TS_{ext} = TS - TS_p$ . We selected events converting in the front section of the tracker, where the Point Spread Function (PSF) is the narrowest. At most, we found  $\sim 3\sigma$  ( $TS \sim 10$ ) evidence for extension (radius of  $0.1^\circ$ ) in the energy range 3–100 GeV. Using all events in the same energy range, the significance of the extension decreases to  $2.5\sigma$ , consistent with the broadening of the PSF. The centroid of the LAT emission is located at 11h12m12s,  $-60^\circ 40' 12''$ , in good agreement with the position of MSH 11–62.

To validate our results we performed a Monte-Carlo simulation of the region, including the diffuse background and the neighboring sources, assuming a Gaussian source of  $0.1^\circ$ . This simulation was analyzed as described above leading to results similar to those obtained with real data. This improves confidence in the  $3\sigma$  evidence for extension. However, more statistics are needed to firmly demonstrate extension of the Fermi-LAT source.

#### 4.2. H.E.S.S. Upper Limits

The position of MSH 11–62 was covered in a survey of the inner Galactic plane with the High Energy Spectroscopic System (H.E.S.S.) in 2004/2005. The approximate exposure time on the source was 9.1 hr (Hoppe 2008), resulting in no reported detection of the source. Based on this non-detection, we calculate a  $2\sigma$  upper limit on the VHE flux at 1 TeV for MSH 11–62 by scaling the H.E.S.S. sensitivity for a  $5\sigma$  point source detection (1% of the Crab in 25 h under the assumption of a photon index of 2.6) to the actual H.E.S.S. exposure for the source. We find  $E^2(dN/dE)_{E=1\text{TeV}} < 3.8 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ , where  $N$  is the number of photons per unit area per second.

## 5. ANALYSIS AND DISCUSSION

### 5.1. Evolutionary State

The large-scale evolution of a composite SNR depends on the mechanical energy of the explosion, the density of the ambient medium, the mass of the supernova ejecta, and the spin-down power of the pulsar. Additional parameters of considerable importance are the braking index and spin-down timescale of the pulsar, along with the fraction of the spin-down power that appears as magnetic flux.

Figure 9 shows an illustrative model for the evolution of MSH 11–62 using parameters for a scenario described in Section 5.3. For these parameters, the forward shock (FS) of the SNR (blue curve) reaches the observed radius of MSH 11–62 at an age of  $\sim 1.2$  kyr. By this time, the RS has begun to approach the PWN, as indicated by the subsequent compression of the PWN radius (red curve). This is consistent with the somewhat distorted morphology of MSH 11–62, which appears to indicate that the RS has propagated most rapidly along the NW/SE direction. In addition, Figure 9 shows that the magnetic field in the PWN has declined to a value of  $\sim 6\mu\text{G}$ .

The broadband emission observed from MSH 11–62 places considerable constraints on the underlying properties of the composite system. Of particular importance is our understanding of the observed  $\gamma$ -ray emission, which could originate from the pulsar, the PWN, or the SNR itself. Here we consider models for each scenario.

### 5.2. Gamma-rays from the SNR

Particle acceleration at the forward shocks of SNRs is known to produce extremely energetic particles that can produce  $\gamma$ -rays through inverse-Compton (IC) scattering of ambient photons by energetic electrons, and/or through the decay of neutral pions produced in energetic




FIG. 10.— Models for the broadband emission from MSH 11–62 under the assumption of dominant hadronic (upper) and leptonic (lower) processes for the observed  $\gamma$ -ray emission.

proton-proton collisions. Thus, one interpretation for the  $\gamma$ -ray emission from MSH 11–62 is that it originates from the SNR itself, and is a signature of particle acceleration by the remnant. We consider two cases, one in which the  $\gamma$ -ray emission is dominated by hadrons ( $\pi^0$ -decay), and one in which it is dominated by leptons (IC scattering).

For the hadronic-dominated scenario, we assume that the mechanical energy of the SNR is  $E_{SNR} = 10^{51}$  erg and that 50% of this energy is converted to energetic particles whose spectrum is described by a power law with an exponential cutoff. We assume an electron-to-proton ratio  $k_{e-p} = 10^{-2}$ , and that the spectral index is the same for protons and electrons. Using  $d = 5$  kpc, we adjust the spectral index, density, and cutoff energy for the proton spectrum to match the Fermi LAT spectrum, while also adjusting the magnetic field strength to reproduce the SNR radio emission with synchrotron emission from the electron spectrum. We find a spectral index of 1.8 and an exponential cutoff energy of 70 GeV for the protons (Figure 10, top), with an ambient density  $n_0 = 6.8 \text{ cm}^{-3}$ . The associated  $\gamma$ -ray emission from IC scattering of the cosmic microwave background (CMB) is negligible. The postshock magnetic field required by the radio emission is  $\sim 10 \mu\text{G}$ . Here we have used the measured 1.4 GHz flux from the shell, and assumed a spectral index range of  $\alpha_r = 0.4 - 0.7$  typical of radio emission from shell-type SNRs.

TABLE 2  
MODEL PARAMETERS:  $\gamma$ -RAYS FROM SNR

Parameter	Hadronic Model	Leptonic Model
$d$	5 kpc	5 kpc
$n_0$	$6.8 \text{ cm}^{-3}$	$0.05 \text{ cm}^{-3}$
$\alpha_e (= \alpha_p)$	1.8	1.8
$E_{cut,e}$	50 GeV	0.9 TeV
$E_{cut,p}$	70 GeV	1 TeV
$b^a$	1.0	2.0
$E_e^b$	$5 \times 10^{48}$ erg	$8 \times 10^{49}$ erg
$E_p^c$	$5 \times 10^{50}$ erg	$8 \times 10^{51}$ erg
$B$	$10.0 \mu\text{G}$	$2.0 \mu\text{G}$

NOTE. —  
a) Curvature index on exponential cut-off  
b) Total electron energy  
c) Total proton energy

The required proton cutoff energy under this scenario is rather low given that we expect energies well in excess of  $10^{12}$  eV to be produced in young SNRs, but it is possible that a low shock speed has resulted in a low maximum energy (e.g., Reynolds 2008). Of potentially greater concern is that the preshock density required under this scenario is more than two orders of magnitude larger than that inferred from the X-ray spectrum. Similar discrepancies have been identified for several  $\gamma$ -ray emitting SNRs associated with OH masers (which imply an interaction between the SNRs and surrounding molecular clouds, thus making  $\pi^0$ -decay a likely production mechanism), and it has been suggested that this could be the result of a clumpy SNR shell in which only the lower-density interclump medium is responsible for the X-ray emission while all of the gas participates in the  $\gamma$ -ray production (Castro & Slane 2010). No maser emission has been reported for MSH 11–62, and the remnant is not known to be interacting with dense material. Thus, while this scenario cannot be ruled out, it does not have strong supporting evidence outside of the  $\gamma$ -ray emission. Searches for evidence of interactions between MSH 11–62 and molecular clouds are of particular importance to further address this picture.

For at least some SNRs, the observed  $\gamma$ -ray emission appears to be dominated by IC emission from an energetic population of electrons (e.g., Ellison et al. 2010). We have considered such a scenario for MSH 11–62 by fixing the ambient density at the value implied by the X-ray measurements, and allowing the normalization and shape of the electron spectrum to vary. We find that for a distance of 5 kpc, the total electron energy required to produce the  $\gamma$ -ray emission through IC scattering of CMB photons is  $8 \times 10^{49}$  erg. The magnetic field required to fit the radio emission from the SNR shell is  $B = 2.0 \mu\text{G}$ . The model, shown in Figure 10 (bottom), requires a super-exponential cutoff of the electron spectrum

$$N_e \propto E_e^{-\alpha_e} \exp\left(-\frac{E_e}{E_{cut,e}}\right)^b \quad (1)$$

Model parameters are given in Table 2. In particular, we find  $E_{cut,e} = 0.9$  TeV, which is quite low given the rather low magnetic field. Since all realistic acceleration models that produce energetic nonthermal electrons in SNRs also




FIG. 11.— Models for the nonthermal emission from MSH 11–62 (lower panel) under the assumption that the  $\gamma$ -ray emission is produced entirely by the PWN. The upper panel shows the associated electron, which is modeled as an evolved Maxwellian distribution with power law tail.

accelerate ions, the total energy budget implied by these results would exceed the available energy of the SNR blast wave for  $k_{e-p} \sim 10^{-2}$  if the remnant distance is 5 kpc.

For the smallest distance allowed by HI absorption measurements, an electron-to-proton ratio  $k_{e-p} > 0.7$  is required for the entire particle energy budget to not exceed  $5 \times 10^{51}$  erg. Such high required values of  $k_{e-p}$ , along with the low cutoff energy and extremely sharp spectral cutoff appear problematic. Thus, we conclude that a leptonic scenario in which the observed  $\gamma$ -ray emission from MSH 11–62 is dominated by IC emission is ruled out for all but the smallest possible distances, and still problematic at such small distance values.

### 5.3. Gamma-rays from the PWN

The X-ray and radio observations of MSH 11–62 clearly identify a powerful PWN in the system. The electrons responsible for the X-ray and radio emission also produce  $\gamma$ -ray emission through inverse Compton scattering. To investigate the scenario in which the Fermi LAT emission is produced entirely by the PWN, we calculate the evolution of the composite system using the model of Gelfand, Slane, and Zhang (2009). We assume expansion of the SNR into a uniform medium of density  $n_0$ , driven by an ejecta mass  $M_{ej}$ . The PWN expansion is powered by the input of a central pulsar with a spin-down power that evolves as

$$\dot{E}(t) = \dot{E}_0 \left[ 1 + \frac{t}{\tau_0} \right]^{-\frac{p+1}{p-1}} \quad (2)$$

where  $\tau_0$  is the initial spin-down timescale for the pulsar, and  $p$  is the braking index. The PWN expansion is confined by the surrounding ejecta. Particles are injected into the nebula with a power law spectrum,

TABLE 3  
MODEL PARAMETERS:  $\gamma$ -RAYS FROM PWN OR PULSAR

Parameter	$\gamma$ -rays from PWN	$\gamma$ -rays from Pulsar
<i>Input:</i>		
$d$	5 kpc	3.5 kpc
$E_{SNR}$	$10^{51}$ erg (fixed)	$10^{51}$ erg (fixed)
$M_{ej}$	$2.5M_{\odot}$	$2.5M_{\odot}$
$n_0$	$0.08 \text{ cm}^{-3}$	$0.03 \text{ cm}^{-3}$
$p$	3.0 (fixed)	3.0 (fixed)
$\tau_0$	1000 yr	1000 yr
$\dot{E}_0$	$4 \times 10^{38} \text{ erg s}^{-1}$	$5 \times 10^{37} \text{ erg s}^{-1}$
$\eta_e^a$	0.999	0.995
$\eta_B^b$	0.001	0.005
$E_{e,min}$	0.1 GeV	1 MeV
$E_{e,max}$	900 TeV	8.5 PeV
$E_p^c$	100 GeV	
$f_{PL}^d$	.1%	
$E_b^e$		440 GeV
$\alpha_e$	2.33	
$\alpha_{e,2}$		2.7
$T_{IR}^f$	25K	25K
$(W_{IR}/W_{CMB})^g$	$2W_{CMB}$	$2W_{CMB}$
<i>Derived:</i>		
Age	1200 yr	720 yr
$B_{PWN}$	$6.2\mu\text{G}$	$13\mu\text{G}$

NOTE. —

- a) Fraction of energy in electrons
- b) Fraction of energy in magnetic field
- c) Peak energy of Maxwellian electron component
- d) Fraction of energy in power law component
- e) Power law break energy
- f) Effective temperature of IR photon field
- g) Ratio of IR energy density relative to CMB

evolved through radiative and adiabatic losses, and the emission spectrum is calculated based upon the evolved magnetic field strength of the nebula at the age required to reproduce the observed radius of the SNR (Figure 9). The ambient radiation field was assumed to be a combination of the cosmic microwave background (CMB) and infrared photons from local dust.

The parameters of the model are summarized in Table 3. We sampled a range of values for  $n_0$  (starting with nominal value determined from the X-ray spectrum in Section 3.4),  $\dot{E}_0$ ,  $\tau_0$ ,  $\alpha_e$ , and  $M_{ej}$ , keeping other parameters fixed at values typical of other pulsars (though we note that parameters such as the braking index and the fraction of the spin-down energy that goes into magnetic flux are not well known). The temperature and energy density of the IR photon field were adjusted slightly to tune the agreement with the data. The final values listed in Table 3 are within reasonable expectations for this region of the Galaxy (see Strong et al. 2000).

We found that it was not possible to reproduce the significant  $\gamma$ -ray emission in the Fermi LAT band with simple power law or broken power law models for the particle spectrum. We then investigated a model consisting of a Maxwellian electron distribution with a power law tail, corresponding to spectra observed in PIC simulations of relativistic shocks (Spitkovsky 2008), considering a range of values for the energy of the Maxwellian peak and for the fraction of the total electron energy found in the power law component. As shown in Figure 11, we



find that the broadband emission from MSH 11–62 can be reproduced to reasonable fidelity for the parameters shown in Table 3. The upper panel of the Figure contains the electron spectrum, and the lower panel presents the synchrotron and IC emission along with the observed data. The radii for the SNR and PWN are shown in Figure 9, along with the PWN magnetic field. The age required to match the observed radius is  $\sim 1.2$  kyr. The magnetic field of the PWN has declined to a value of  $\sim 6\mu\text{G}$ , and the PWN radius is approaching the stage at which it encounters the SNR reverse shock, as indicated by the subsequent decline in  $R_{PWN}$ . This appears consistent with the X-ray and radio morphology, as discussed above, although the observations seem to suggest that the interaction has proceeded beyond that indicated in Figure 9. We note that the magnetic field in this scenario is far below the equipartition value calculated from radio measurements.

According to this model,  $\gamma$ -ray emission in the Fermi LAT band is produced entirely by electrons in the Maxwellian peak of the spectrum. This is consistent with results for Vela X (de Jager, Slane, & LaMassa 2008), HESS J1640–465 (Slane et al. 2010), and several other PWNe (Fang & Zhang 2010) that suggest a low-energy electron component distinct from the power law component. For MSH 11–62, we find a mean value of  $\gamma = 2 \times 10^5$  for the electrons in the Maxwellian component, with only 0.1% of the total energy in the power law tail.

We note that the initial spin-down power required for this model,  $\dot{E}_0 = 4 \times 10^{39}$  erg s $^{-1}$ , is uncomfortably large. For the assumed braking index  $p = 3$ , this would lead to a value of  $\dot{E} = 8.3 \times 10^{38}$  erg s $^{-1}$  at the current epoch – higher than that for any known pulsar. This would imply  $L_x/\dot{E} \sim 10^{-6}$  for the pulsar, which is extremely low.

For a distance of 2 kpc, reasonable fits to the broadband spectrum can also be obtained, with a current spin-down power of  $\sim 2 \times 10^{38}$  erg s $^{-1}$ . While still extremely high, this is similar to the value for PSR J2022+3842, a pulsar recently discovered in G76.9+1.0 (Arzoumanian et al. 2011). This pulsar is characterized by an extremely low conversion efficiency of spin-down power into X-rays as well, but the value ( $L_x/\dot{E} \sim 6 \times 10^{-5}$ ) is still much higher than that inferred for the pulsar in MSH 11–62 if the Fermi LAT emission is being produced by the PWN. We conclude that it is possible for the observed  $\gamma$ -ray emission from MSH 11–62 to be produced by the PWN, but that this would require a very significant amount of energy in the Maxwellian peak of the electron spectrum, leading to spin-down parameters for the pulsar that seem unlikely.

#### 5.4. Gamma-rays from the Pulsar

Fermi observations have clearly established that pulsars comprise a significant fraction of the Galactic population of GeV point sources. The associated spectra are characterized by a power law with an exponential cutoff, with  $E_{\text{cut}} \sim 1 - 6$  GeV, and  $\Gamma \sim 1 - 2$  (Abdo et al. 2010). This is very similar to the spectrum observed for MSH 11–62. Being associated with a composite SNR, for which our X-ray observations identify the underlying neutron star, it is quite possible that the observed  $\gamma$ -ray

emission originates with the pulsar itself.

The ratio of the X-ray flux from the point source in MSH 11–62 to the observed  $\gamma$ -ray flux is  $F_x/F_\gamma \sim 5 \times 10^{-3}$ . This value is typical of that for pulsars detected by the Fermi LAT, although there is a large spread in this ratio for the population.

Using the approximate relationship  $L_x \approx 10^{-3}\dot{E}$  derived from a variety of studies of X-ray PWNe and their associated pulsars, the spectrum of the PWN in MSH 11–62 suggests an underlying pulsar with  $\dot{E} \sim 10^{37}d_5^2$  erg s $^{-1}$ . We note, however, that there are large variations in the observed relation between  $L_x$  and  $\dot{E}$  (e.g., Vink, Bamba, & Yamazaki 2011). More than 85% of the pulsars detected by the Fermi LAT, and all of those pulsars discovered in the  $\gamma$ -ray band, have lower values of  $\dot{E}$ . However, only one pulsar discovered in  $\gamma$ -rays has a Fermi LAT flux lower than that associated with MSH 11–62. Thus, the lack of identified pulsations in any band is not strong evidence that the emission from MSH 11–62 is not associated with the pulsar itself. The marginal evidence for extension of the  $\gamma$ -ray emission, which would rule out the pulsar as the source of the emission, is of particular importance in this context, and requires revisiting in the future when additional data from this source have been obtained.

If we assume that the Fermi LAT emission arises entirely from the pulsar in MSH 11–62, the PWN emission can then be described by a broken power law model with  $\dot{E}_0 \sim 5 \times 10^{37}$  erg s $^{-1}$  and an age of  $\sim 700$  yr, at which time an ambient density  $n_0 = 0.03$  cm $^{-3}$  yields the observed radius for the SNR. The resulting model spectrum is shown in Figure 12, and the parameter values for the model are listed in Table 3. The PWN magnetic field strength is  $\sim 13\mu\text{G}$ , a factor of five lower than the equipartition field derived from radio observations, and the spin-down power at the current epoch is  $\dot{E} = 1.7 \times 10^{37}$  erg s $^{-1}$ . This leads to  $L_x/\dot{E} \approx 3 \times 10^{-4}$ , well within the observed range for known pulsars. In this model, the SNR RS has not yet begun to impact the PWN, leaving the observed X-ray and radio morphology unexplained.

## 6. CONCLUSIONS

Radio observations of MSH 11–62 indicate that this is a composite SNR containing a central PWN. X-ray observations confirm this, with clear identifications of the PWN synchrotron emission as well as the presence of an X-ray point source that most likely corresponds to the pulsar driving the system. The X-ray spectral index in the PWN increases with distance from the compact source, as expected from synchrotron losses, and the strongly asymmetric morphology of the PWN suggests an interaction with the SNR RS.

Observations with the Fermi LAT reveal distinct  $\gamma$ -ray emission from MSH 11–62, but the limited angular resolution makes it impossible to ascertain the origin of the emission directly. We have considered three cases, with the  $\gamma$ -ray emission arising from the SNR, the PWN, and the pulsar, respectively. Models for each scenario can reproduce basic features of the observed broadband nonthermal emission, but not without difficulties. If the emission arises from the SNR, then a hadronic scenario requires an ambient density that is a factor of




FIG. 12.— Model for the PWN emission from MSH 11–62 under the assumption that the  $\gamma$ -ray emission arises from the pulsar. Parameters for the model are given in Table 2.

$\sim 100$  higher than that implied by the thermal X-ray emission, while a leptonic scenario requires that an unreasonably large fraction of the SNR mechanical energy is converted to relativistic particles unless the SNR distance is quite low. Both models require very low energy cutoffs in the particle spectra. If the emission arises from the PWN, then the input spectrum from the pulsar requires a dominant Maxwellian component, with only 0.1% of the energy in the accompanying power law tail, and an extremely high pulsar spin-down power that appears inconsistent with the observed X-ray and  $\gamma$ -ray flux values.

The most likely scenario is that the  $\gamma$ -ray emission arises from the pulsar itself. While models for the PWN emission still imply a somewhat high spin-down power, the lack of detected  $\gamma$ -ray pulsations is not surprising given the relatively low flux of the source. The weak evidence for extent in the  $\gamma$ -ray emission is not sufficient to reject this scenario, but is of considerable interest for additional study with future observations.

NASA supported this work via grant numbers NRA 00-OSS-07/03500279 and NGT5-159. PS acknowledges support from NASA Contract NAS8-03060. The ATCA is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO.

#### REFERENCES

- Abdo, A. A., et al. 2009, ApJ, 706, 1331  
 Abdo, A. A., et al. 2010, ApJS, 187, 460  
 Abdo, A. A. et al. 2011, arXiv:1108.1435v1  
 Arzoumanian, Z., et al. 2011, arXiv:1105.3185  
 Castro, D. & Slane, P. 2010, ApJ, submitted (arXiv:1002.2738)  
 de Jager, O., Slane, P., & LaMassa, S. 2008, ApJ, 689, L125  
 Ellison, D. C., et al. 2010, ApJ, 712, 287  
 Fang, J. & Zhang, L. 2010, A&A, 515, A20  
 Gaensler, B. M. & Slane, P. 2006, ARAA, 44, 17  
 Gelfand, J.D. et al. 2007, ApJ, 663, 468  
 Gelfand, J. D., Slane, P. O., & Zhang, W. 2009, ApJ, 703, 2051  
 Grondin, M.-H., et al. 2011, ApJ, 738, 42  
 Harrus, I. M., Hughes, J. P., & Slane, P. 1998, ApJ, 499, 273  
 Harrus, I. M., et al. 2002, Neutron Stars in Supernova Remnants, ASP Conference Series, Vol. 271, held in Boston, MA, USA, 14-17 August 2001. Edited by Patrick O. Slane and Bryan M. Gaensler. San Francisco: ASP, 2002., p.221  
 Hartman, R.C. et al. 1999, ApJS, 123, 79  
 Hoppe, S. 2008, Proceedings of the 30th International Cosmic Ray Conference Rogelio Caballero, Juan Carlos D’Olivo, Gustavo Medina-Tanco, Lukas Nellen, Federico A. Sanchez, Jos F. Valds-Galicia (eds.) Universidad Nacional Autnoma de Mexico, Mexico City, Mexico, 2008 Vol. 2, pages 579-582  
 Johnston, S., McCure-Griffiths, N. M., & Koribalski, B. 2004, MNRAS, 348, L19  
 Kargaltsev, O. & Pavlov, G. 2008, AIP 983, 171  
 Kaspi, V. M., et al. 1997, ApJ, 485, 820  
 Kennel, C. F., & Coroniti, F. V. 1984, ApJ, 283, 694  
 Lazendic, J. S., et al. 2000, ApJ, 540, 808  
 Mattox, J. R., et al. 1996, ApJ, 461, 396  
 Moffett, D., Gaensler, B.M.; Green, A. 2001, Young Supernova Remnants: Eleventh Astrophysics Conference. AIP Conference Proceedings, Volume 565, pp. 333-336 (2001)  
 Moffett, D., Gaensler, B.M., Green, A., Slane, P., Harrus, I.M., Dodson, R. 2002, Neutron Stars in Supernova Remnants, ASP Conference Series, Vol. 271, held in Boston, MA, USA, 14-17 August 2001. Edited by Patrick O. Slane and Bryan M. Gaensler. San Francisco: ASP, 2002., p.221  
 Reynolds, S. P. 2003, Proceedings of IAU Colloquium 192, 10 Years of SN1993J (Valencia, Spain, April 2003) Edited by J.M. Marcaide & K.W. Weiler Springer-Verlag  
 Reynolds, S. P. 2008, ARA&A, 46, 89  
 Roger, R. S., Milne, D. K., Caswell, J. L. and Little, A. G. 1986, MNRAS, 219, 815  
 Slane, P., et al. 2000, ApJ, 533, L295  
 Slane, P., et al. 2004, 616, 403  
 Slane, P. et al. 2010, ApJ, 720, 266  
 Spitkovsky, A. 2008, ApJ, 682, L5  
 Strong, A. W., Moskalenko, I. V., & Reimer, O. 2000, ApJ, 537, 763  
 Sturmer, S. J. & Dermer, C. D. 1995, A&A, 293, L17  
 Tam, C., Roberts, M. S. E., & Kaspi, V. M. 2002, ApJ, 572, 202  
 Vazquez, R. A. & Feinstein, A. 1990, A&AS, 86, 209  
 Vink, J., Bamba, A., & Yamazaki, R., 2011, ApJ, 727, 131  
 Wilson, A. S. 1986, ApJ, 302, 718