





# Dark Matter


- Dark Matter is there but **we know nothing about the particle content of DM**
  - Plenty of cosmological/ astrophysical observations: CMB anisotropies, galaxy rotation curves, gravitational lensing, cluster collisions...
- No hints on DM particle properties (mass, cross section)
- Common assumption: **thermal origin of DM:**
  - DM in thermal equilibrium with SM in early Universe. Current relic abundance set by the strength of the SM-LDM interaction (“freeze-out mechanism”)
  - constrain on available mass range
- Light Dark Matter: mass range  $1 \text{ MeV} \div 1 \text{ GeV}$






# Dark Photon Production Mechanisms With Lepton Beams

Three main production mechanisms in fixed targets, lepton beam experiments:




## a) $A'$ -strahlung:

- Radiative  $A$  emission in nucleus EM field
- Scales as  $Z^2\alpha_{EM}^3$ .
- Forward-boosted, high-energy  $A$  emission



## b) $e^+e^-$ annihilation:


- scales as  $Z\alpha_{EM}^2$ .
- Forward-backward  $A'$  emission in the CM



## c) Resonant $e^+e^-$ annihilation

- scales as  $Z\alpha_{EM}$ .
- resonant Breit-Wigner like cross section with  $m_{A'} = \sqrt{2m_e E}$




# Mass Hierarchy Determines Search Strategy and Interpretation



- (a) Secluded scenario: does not lend itself to decisive laboratory tests.
- (b) Visible decay scenario.  
Experiments @ JLAB: HPS, APEX
- (c) Invisible decay scenario.  
Experiments @ JLAB: BDX, BDX-MINI

# APEX: A-Prime EXperiment

- e- fixed target experiment installed in **HALL A**.
- Dark photon searched as a narrow resonance in  $e+e-$  mass over a smooth QED background
- Two High Resolution Spectrometers (HRSs) in coincidence to measure events with an  $e-$  in one arm and  $e+$  in the other
  - Standard HRS detector stack in both arms: Scintillators: S0 and S2(timing), VDC (tracking), Cherenkov and Calorimeters (PID)




- 2010 test run<sup>a</sup>: beam 2.2 GeV@150 uA on tantalum foil
- Full run in 2019: accumulated over 100x statistics than test run

<sup>a</sup> 10.1103/PhysRevLett.107.191804

# HPS: Heavy Photon Search

e<sup>-</sup> fixed target experiment installed in **HALL B**.



## Key points

- CEBAF e<sup>-</sup> beam: 1.1-6.6 GeV
- Thin W target :  $10^{-3} \times 0$
- Si tracker inside a dipole magnet: momentum measurement / vertexing
- PbWO<sub>4</sub> calorimeter <sup>a</sup> downstream: PID / trigger


## Signature

- Resonance search (aka “bump-hunt”): Narrow e<sup>+</sup>e<sup>-</sup>-resonance over a QED background ( $\epsilon \sim 10^{-3}$ )
- Detached vertex search: Search for two tracks showing a common production vertex downstream the target ( $\epsilon \sim 10^{-4}$ ).

<sup>a</sup> I. Balossino et al., NIMA 854, (2017) 89

# HPS Status and Results

- HPS already completed 2 “engineering” runs (2015<sup>a</sup>/2016<sup>b</sup>) and 2 “production” runs (2019/2021).
- Results from engineering runs allowed to optimize the detector and demonstrate the HPS capabilities - upper limits set for the  $A'$  parameters space, although no new regions were explored.
- Results from production runs will investigate for the first time unexplored territories in the  $A'$  space - analysis ongoing, stay tuned for results!
- More runs to come! 102 “PAC” measurement days ( 204 calendar days) still to run.



<sup>a</sup> P.H.Adrian et al. (Heavy Photon Search Collaboration), Phys. Rev. D 98, 091101(R)

<sup>b</sup> arxiv:2212.10629 - accepted by PRD





# X17 search at JLab

Direct detection search for hidden sector particles using the magnetic-spectrometer-free PRad setup in **Hall-B**<sup>a</sup>:



- Discover or establish an experimental upper limit on the electroproduction of the hypothetical X17 particle, claimed in two ATOMKI low-energy proton-nucleus experiments.
- Search for “hidden sector” intermediate particles in [3–60] MeV mass range
- Experimental approach:
  - bump hunt
  - direct detection of all final state particles ( $e^+$ ,  $e^+e^-$  or  $\gamma\gamma$ )

<sup>a</sup> arXiv:2108.13276v3



# BDX: Beam Dump eXperiment

- BDX is a JLAB experiment approved by PAC46
- Detector installed O(20 m) behind **Hall-A beam-dump**
- Two step experiment:
  - **production** of LDM beam
  - **detection** of LDM particle: scatters on  $e^-$  in the detector realising visible signal (O(GeV) EM shower)




## Key points

- 11 GeV  $e^-$  beam, current: 65  $\mu\text{A}$
- charge:  $1\text{E}22$  EOT
- Fully parasitic wrt Hall-A physics program
- New experimental hall behind Hall A beam dump
- BDX detector:
  - EM calorimeter: CsI(Tl) crystals+SiPM readout (active volume  $\sim 0.5\text{ m}^3$ )
  - Dual active-veto layer made of plastic scintillator counters + SiPM readout
  - Passive lead layer surrounding the calorimeter

## BDX sensitivity

BDX will improve of 2 orders of magnitude current exclusion limits in LDM parameter space





# BDX-MINI@JLAB: Pilot experiment

Small-scale, low energy version of full BDX experiment

## Experimental setup

- 2.2 GeV, 150  $\mu\text{A}$   $e^-$  beam
- SM particles shielded by concrete and soil
- Detector installed in a well 25 m downstream



## BDX-MINI detector<sup>a</sup>


- PbWO<sub>4</sub>-based EM calorimeter (44 crystals), SiPM readout
  - 0.15% of BDX active volume
- 8 mm passive Tungsten shielding
- 2 plastic scintillator active veto layers, SiPM readout

<sup>a</sup> M. Battaglieri et al., EPJ C 81(2021)164

## BDX-MINI@JLAB: Results

- **Blind approach:** fix the selection cuts by optimizing the experiment sensitivity.
- Upper limit is derived in the LDM parameters space
- This pilot experiment<sup>a</sup> is sensitive to the parameter space covered by some of the most sensitive experiments to date.


<sup>a</sup> M.Battaglieri and et al, Phys. Rev. D 106, 072011



## BDX @ 22 GeV

- BDX@22 GeV<sup>a</sup> can complement BDX measure
- 90% CL BDX sensitivity:
  - Ideal case of a zero-background measurement
  - energy threshold: 300 MeV
  - an overall 20% signal efficiency.



<sup>a</sup> arXiv:2306.09360



# Probing muon-philic forces with secondary muon beam

- Sizable flux of high energy muons produced in a thick target by the interaction of the e-beam:
  - 11 GeV (22 GeV) e- hitting Hall-A beam dump:  $3 \cdot 10^8 \mu/s$  ( $\sim 2 \cdot 10^9 \mu/s$ )
  - expect muon flux @ KL-facility:
- Use the secondary muon beam to produce exotic particles accounting for g-2 anomaly

$\mu$ BDX<sup>2</sup>: Muon beam dump experiment to probe the visible decay into  $e^+e^- (\gamma\gamma)$




<sup>2</sup> L. Marsicano et al., PRD 98, 115022 (2018)

# Probing muon-philic forces with secondary muon beam

- Sizable flux of high energy muons produced in a thick target by the interaction of the e-beam:
  - 11 GeV (22 GeV) e- hitting Hall-A beam dump:  $3 \cdot 10^8 \mu/s$  ( $\sim 2 \cdot 10^9 \mu/s$ )
- Use the secondary muon beam to produce exotic particles accounting for g-2 anomaly


Fixed-target, missing-momentum search strategy to probe invisibly decaying particles (M3 experiment @Fermilab like)






# LDM search using $e^+$ beam: Missing energy approach

- Missing energy<sup>a</sup> experiment with a 11 GeV positron beam
- $e^+$  impinging on active thick target (ECAL);  $A'$  produced via resonant process  $e^+e^- \rightarrow A'$
- large missing energy as LDM production signature:  $E_{miss} = E_{beam} - E_{ECAL}$
- HCAL to detect neutral particles escaping the ECAL mimicking signal



<sup>a</sup> M. Battaglieri et al., Eur. Phys. J. A 57, 253 (2021)



# Conclusions

- Jefferson Lab features a rich LDM experimental program (HPS, BDX-mini, APEX)
- New developments are expected in the nearby future: the Beam Dump eXperiment can run in the next few years provided the new hall is built
- The realization of a positron beam at Jefferson Lab paves the way to new competitive LDM experiments
- High-intensity muon secondary beam will complement the current program