



Understanding Potential Instrumental Effects in Searches for Narrow Spectral Features



Eric Charles, A. Albert

**E. Bloom, A. Snyder, Z. Yang,
and G. Zaharijas**

**on behalf of the Fermi-LAT
collaboration**

**4th Fermi Symposium
Monterey CA**

The Context: Narrow Feature at 130 GeV




Bringmann et al. and Weniger showed evidence for a narrow spectral feature near 130 GeV near the Galactic center (GC).

- Signal is particularly strong in 2 out of 5 test regions, shown above.
- Over 4σ , with S/N > 30%, up to ~60% in optimized regions of interest (ROI).
- Some indication of double line (111 & 130 GeV).


What the LAT Team Sees

Signal significance in fit to powerlaw + 1D signal PDF at 130 GeV for $4^\circ \times 4^\circ$ boxes near the Galactic Center in 1° steps.



Significant ($3-4\sigma+$) excess near the Galactic center, slight offset towards negative latitudes. Largely within 4° of Galactic plane. (However, be wary of interpretation in view of limited statistics...)


Slicing Along the Plane



At 130GeV the excess near the Galactic Center might have structure. (Again, be wary of interpretation in view of limited statistics). Feature at -10° maybe be noise, but it is almost the same size as the feature at the GC.

Signal significance in fit to powerlaw + 1D signal PDF at 130GeV for $4^\circ \times 4^\circ$ boxes along the galactic plane.

Morphology of “Two-Line” Excesses



The 111GeV feature is not as strong as the 130GeV feature, and does not appear to be spatially correlated with it, expect perhaps in the $4^\circ \times 4^\circ$ ROI nearest the Galactic center.

Systematic Effect to Consider

Effects	Scaling	Visible with peak to off-peak event comparison?
Particle Backgrounds	Solid Angle	Yes
Increased A_{eff} at E_{peak}	γ rays in Energy Band	Probably
Decreased A_{eff} near E_{peak}	γ rays in Energy Band	Maybe
Energy Redistribution	γ rays in Energy Band	Maybe

It is difficult to disentangle the three final cases when only looking at events in the final sample.


Decreased A_{eff} near E_{peak} is particularly challenging, as the events of interest are not in the final sample.

We have also considered many other effects: e.g., point source masking, simulated effects not captured in IRFs, data/ MC comparisons of selection variables, etc...

See also: [[arXiv:1209.4548](#)], [[arXiv:1209.4562](#)].

INSTRUMENTAL CONSIDERATIONS IN STUDYING NARROW SPECTRAL FEATURES


Observing Profiles Variations



Averaged over years, the observing profile depends primarily on the DEC of the Region of Interest (ROI).

The Galactic Center gets somewhat more time right on-axis than other sources (and less time slightly off-axis). This is because $\text{DEC}_{\text{GC}} \sim \text{Inclination}_{\text{orbit}}$

θ -averaged Energy Resolution by Declination




The θ -averaged PDF weighted for observing profile varies moderately with declination.

Using the wrong profile will not induce a signal, but can scale the significance of a signal by up 25%. This suggests using a 2 dimensional PDF, which increases statistical power by $\sim 15\%$ (see Andrea Albert's talk).

BACKGROUND CONTAMINATION


Cosmic-ray Background Contamination



Above 100 GeV most of the high-latitude events in P7SOURCE that are not in P7CLEAN are not γ rays.

CR-background reconstructed as γ rays will show a variety of spectral features, which can corrupt and compromise the sideband fit as well as induce fake signals. (See E. Bloom talk).

Background Contamination Rates




ApJS, 203, 4. [[arXiv:1206.1896](https://arxiv.org/abs/1206.1896)]

Comparing P7SOURCE fits for small ROI in the galactic plane to large ROI where the P7SOURCE class is dominated by CR background is dangerous.

For very large ROI the residual contamination in the P7CLEAN class may becoming noticeable, but this is **not** an issue for small ROI near the GC, and is **not** causing the feature seen at 130GeV.


EFFICIENCY AS A FUNCTION OF ENERGY

Measuring Efficiency with the Earth Limb



The Earth Limb is unique in that it can be seen in the P7TRANSIENT class at high energies. This allows us to use it to measure efficiencies for tighter event classes as a function of energy.


P7TRANSIENT to P7CLEAN Efficiency



These dips in efficiency appear to be related to the CAL-TRK agreement.


The efficiency at $\sim 115 \text{ GeV}$ is $0.57/0.75 = 75\%$ of the MC prediction. This would imply a 30% boost in signal at 130 GeV relative to the prediction from nearby energy bins.

Fitting the Earth Limb



The fit to Earth Limb data results in a 3.0σ signal, with a fractional residual (i.e., S/N) of $\sim 18\%$.



Earth Limb Residuals as a Function of Energy



Fits to Earth Limb data at $\sim 1.5\%$ energy steps

The limited statistics at high energies make it difficult to exclude large fractional residuals. At 130GeV we see $S/N \sim 18\%$ and 3.0σ significance residuals.

Signal to Noise of Excess as a function of θ





Many people have noted that the spectral excess in both the the GC and the earth limb is largest near $\cos(\theta)=0.7$.

By comparing the fractional residuals we see that the features in the Earth Limb could account for about 50% of the excess in a 12°x12° box around the GC, but only about a 30% of the excess in a smaller 8°x8° box where the feature is brighter.

NEW AND UPCOMING DEVELOPMENTS


Data Reprocessing with Updated Calibrations



Reprocessing Data with updated calibrations (primarily CAL). See poster by J. Bregeon. Improves the CAL/TKR agreement at high energies, improving the PSF. Also corrects for loss in CAL light yield b/c of radiation damage (~4% in mission to date) which corresponds to a slightly larger change in the energy scale at 130 GeV. 80%+ overlap in events between original and reprocessed samples.

Results largely similar. Spectral feature has shifted to 135 GeV and the significance has decreased slightly. See talks by E. Bloom and A. Albert. Figures with P7REPCLEAN attached to this talk.

Pass 8 Event Reconstruction



Improved TKR and CAL reconstruction algorithms mitigate issues with CAL /TKR agreement, help avoid features in A_{eff} curve. (See poster by L. Baldini)

Expect ~25% increase in acceptance above ~10 GeV from using improved reconstruction information for event selection (See poster by M. Wood)

Expect better energy resolution at high energies from improved shower profile fitting (See poster by P. Bruel)



Pass 8 event analysis, nearing completion and expected in 2013 will substantially improve our prospects for answering questions about the spectral feature at 130 GeV.

Summary

- **Spectral feature at 130 GeV near the Galactic center is a potentially interesting hint of Dark Matter annihilation**
 - **Fractional residual up to 60% in 4°x4° box around GC**
 - **Not caused by background contamination**
- **A similar spectral feature is seen in the Earth Limb and is likely attributable to dips in efficiency at energies just above and below 130 GeV**
 - **The Earth Limb instrumental features are not enough to explain all of the feature near the GC, however when accounted for they reduce the significance of the GC feature by up to 30%-50% depending on the ROI under consideration.**
- **Data have been reprocessed with updated CAL calibrations**
 - **Detailed results to be presented in talks by E. Bloom and A. Albert**

EXTRA SLIDES


Performance Variation with Incident Angle (θ)



Because of the shape of the LAT, the γ -ray incident angle θ is the key parameter for performance.

Changing resolution with θ implies that cutting on a Energy band to identify signal events will shape the distribution in θ and increase the “mistag” rate of between signal and background. See [[arXiv:1208.3677](https://arxiv.org/abs/1208.3677)] for a better method.


Where are we most sensitive to a narrow feature



Out to about $\theta=50^\circ$, the improving energy resolution balances out the decreasing A_{eff} . Less sensitivity past $\theta=60^\circ$.

Toy MC simulations for a range of signal-to-noise ratios favor energy resolution over A_{eff} slightly less than naïve scaling predictions.


Zenith Cuts Shape the Observing Profile




Standard zenith angle selection reduces the amount of observing time away from the boresight because the LAT tends to point away from the Earth.

“Effective Background”

Signal and Bkg. PDFs



Signal Weight v. Energy



Signal Weight: $W_s(E) = P_s(E) / (P_s(E) + P_b(E))$

“Effective Bkg”: $N_{\text{bkg,eff}} = N_{\text{evt}} \int P_b(E) W_s(E) dE$

Significance: $S = N_{\text{sig}} / \text{sqrt}(N_{\text{bkg,eff}})$

Fractional Residual(i.e., S/N): $F = N_{\text{sig}} / N_{\text{bkg,eff}}$

To consider instruments effects it is useful to look at the potentially induced fractional residual (i.e., the Signal-to-noise ratio).


Event Selection Cuts

Selection	CUT	Comments
P7TRANSIENT	Quality Cuts	
	Charged Particle Veto Analysis	
	Loose cut on P_{all} (0.2)	Small feature in MC (S/N ~ 0.05)
P7SOURCE	CAL & TKR Vetos	
	Reject MIPs with CAL & TKR	
	CAL / TKR Agreement	
	PSF Quality	Depends on CAL/TKR agreement
	Tight cut on P_{all} (0.996 at 130GeV)	Depends on CAL/TKR agreement
P7CLEAN	Reject MIPs, but lose A_{eff}	
	Shape of event in CAL	
P7ULTRACLEAN	Tighter cut on P_{all} below 10GeV	


The two cuts in red appear to account for most of the difference between Earth Limb data and MC at high energies.

Acceptance of Event Classes

Acceptance v. Energy



Events Rates over a Day




ApJS, 203, 4. [[arXiv:1206.1896](https://arxiv.org/abs/1206.1896)]

Above a few GeV: P7TRANSIENT class primarily rejects poorly reconstructed events and events which fail the ACD analysis due to backslash from the CAL with very loose cuts on CAL and TKR topology.

P7SOURCE, P7CLEAN and P7ULTRACLEAN class make more use of the TKR and CAL and reject MIP-like events, events that look more like hadronic showers, and events with poor CAL/TKR agreement.


Observing Profile of the Earth Limb



The observing profile for pointed-mode data, with the standard rocking angle cut reversed: ($\text{ABS}(\text{ROCK_ANGLE}) > 52$) is similar to the GC for larger off-axis angles, but under-represents angles near the boresight.

Note a scaling factor of > 400 between the observing times in the two samples.

CAL/TKR Agreement, High Energy PSF, etc..





CalTrackAngle: angle between CAL axis and TKR direction

CalTrackDoca: Distance of Closest Approach (DOCA) between track and CAL centroid

P_{CORE} : Probability that event is within the CORE of the PSF

Above $\sim 10\text{GeV}$ the backslash from the CAL causes many hits in the TKR and increases the probability of picking the wrong hit for a track and pulling the track direction well into the tails of the PSF. We use the TKR / CAL agreement to mitigate this and also to reduce CR background.

Data / Monte Carlo Comparisons




Detailed comparisons between flight data and Monte Carlo simulations show that the CAL/TKR agreement is somewhat worse in the flight data than in the simulations.

These two variables are among the most important in the Classification Tree analyses used for event selection and classification.

THE GALACTIC PLANE AS A TEST SAMPLE


Fitted Excesses Along the Plane at 89GeV



The significance distributions look like noise with a few noticeable features of marginal significance (e.g., $l, b = -28^\circ, -4^\circ$).

Signal significance in fit to powerlaw + 1D signal PDF at 130 GeV for $4^\circ \times 4^\circ$ boxes near along the Galactic plane in 1° steps.


Fitted Excesses Along the Plane at 130 GeV



The excesses near the GC at 130 GeV are among the largest seen at any energy at the $4^\circ \times 4^\circ$ scale, and stand out particularly at 8-16° scales.

Integrating the Galactic plane outside $\pm 10^\circ$ shows no excess at 130 GeV. (See Elliott Bloom's talk).

Distribution of Significances




Distribution of significances for all ROI centered within a 20°x40° around the Galactic center. Signal energy scanned in 0.25 σ steps between 65 and 500 GeV and $\pm 6\sigma$ fit windows. Fits are to a powerlaw + 1D signal PDF.

The red histogram shows fits using data with shuffled energies. The black histogram is flight data. The blue curve is a fit to the shuffled data. In both cases it is consistent with a unit width Gaussian with mean zero.

As stated earlier, the excess at 130GeV is among the largest seen at any energies, and stands out from the distribution at 8° to 16° scales.


LAUNCH AND EARLY ORBIT LIMB DATA

CAL Temperatures During Early Orbit




Commissioning data used slightly different configurations, and original processing had poor CAL intra-range calibrations.

CAL Temperatures During Survey Mode



RESULTS WITH REPROCESSED DATA


Slice Along the Galactic Plane



In the reprocessed data the hints of sub-structure are more pronounced. $P(\chi^2)$ for central 5(7) bins to be flat is 2% (2×10^{-4}).


Signal significance in fit to powerlaw + 1D signal PDF at 136GeV for $4^\circ \times 4^\circ$ boxes along the galactic plane.

Fitted Excesses Along the Plane at 93GeV



Fit significance for $4^\circ \times 4^\circ$ ROI in 1° Steps along the Galactic Plane at 93 GeV.
(This corresponds to 89 GeV in the original data set).

Fitted Excesses Along the Plane at 136GeV



Fit significance for $4^\circ \times 4^\circ$ ROI in 1° Steps along the Galactic Plane at 136 GeV.
(This corresponds to 130 GeV in the original data set).