



*Marscher et al. 2008, Nature 452, 966*



*Kikuchi et al., 1980, A&A, 190, L8*



*Abdo et al. 2010, Nature 463, 919*

# RoboPol: the optical polarisation of a $\gamma$ -ray flux limited sample of AGN

Emmanouil Angelakis<sup>1</sup>

D. Blinov<sup>2,3</sup>, V. Pavlidou<sup>2,3</sup>, T. Hovatta<sup>4</sup>, I. Myserlis & the RoboPol collaboration

<sup>1</sup>Max-Planck-Institut fuer Radioastronomie, Auf dem Huegel 69, 53121 Bonn, Germany

<sup>2</sup>Foundation for Research and Technology - Hellas, IESL, Voutes, 7110 Heraklion, Greece

<sup>3</sup>Department of Physics and Institute for Plasma Physics, University of Crete, 71003, Heraklion, Greece

<sup>4</sup>Aalto University Metsahovi Radio Observatory, Metsahovintie 114, 02540 Kylmala, Finland



MAX-PLANCK-GESELLSCHAFT



Max-Planck-Institut  
für Radioastronomie


# the RoboPol program


*Pavlidou, EA et al. 2014, MNRAS, 442, 1693*

- ➔ unbiased samples:
  - ▶ 65 GL sources: from 2FGL
  - ▶ 15 GQ sources: variable in radio
- ➔ adaptive cadence: 3 - 0.3 nights
- ➔ 4-channel RoboPol polarimeter  
*King et al. 2014, MNRAS, 442, 1706*  
*Ramaprakesh et al., in prep.*



**Caltech:** M. Balokovic, A. Mahabal, T. J. Pearson, A. Readhead  
**Uni of Crete:** D. Blinov, N. Kylafis, G. Panopoulou, I. Papadakis, I. Papamastorakis, V. Pavlidou, P. Reig, K. Tassis  
**MPIfR:** E. Angelakis, I. Myserlis, J. A. Zensus  
**IUCAA:** V. Joshi, S. Prabhudesai, A. Ramaprakash  
**Nicolaus Copernicus University:** A. Kus - A. Marecki, E. Pazderski  
**Other:** T. Hovatta, S. Kiehlmann, O. King






season 2013



season 2014






season 2013



season 2014

- ➔  $p$  uncertainty: less than 0.01
- ➔  $\chi$  uncertainty: 1-2 deg
- ➔  $R$ -mag uncertainty:  $\sim 0.02$ - $0.04$  mag



median (KS test p:  $6.5 \times 10^{-4}$ )

➔ GL: 0.078

➔ GQ: 0.031

$$\text{PDF} = \frac{1}{x\sigma\sqrt{2\pi}} e^{-\frac{(\ln x - \mu)^2}{2\sigma^2}}$$


➔ GL: 0.105 (var: 0.0068)

➔ GQ: 0.035 (var: 0.0011)

the polarization of GL and GQ:

*Angelakis et al. in prep.*




- GL more polarized than GQ:
  - ▶ uniformity of the field?
- function of the synchrotron peak



# the polarization of GL and GQ:

*Angelakis et al. in prep.*

- GL more polarized than GQ:
  - ▶ uniformity of the field?
- function of the synchrotron peak
- independent of luminosity:
  - ▶ no association with source class




*Angelakis et al. in prep.*

## the polarization of GL and GQ:

*Angelakis et al. in prep.*

- ➔ GL more polarized than GQ:
    - ▶ **uniformity of the field?**
  - ➔ function of the synchrotron peak
  - ➔ independent of luminosity:
    - ▶ no association with source class
  - ➔ independent of the radio variability amplitude
- Richards et al., 2011, ApJS, 194, 29*
- ➔ correlated with the optical variability amplitude
  - ➔ non-thermal events?
  - ➔ a mechanism that:
    - ▶ moves the SED horizontally
    - ▶ increases the polarisation



*Angelakis et al. in prep.*




# EVPA rotations

*Blinov et al. 2015, MNRAS.453.1669B; Blinov et al. in prep.*

→ detected 27 rotations:

- ▶ 2013: 16 rotations in 13 blazars  
*Blinov et al. 2015, MNRAS.453.1669B*
- ▶ 2014: 11 rotations in 10 blazars  
*Blinov et al. in prep.*



# EVPA rotations

*Blinov et al. 2015, MNRAS.453.1669B; Blinov et al. in prep.*

→ detected 27 rotations:

▶ 2013: 16 rotations in 13 blazars

*Blinov et al. 2015, MNRAS.453.1669B*

▶ 2014: 11 rotations in 10 blazars


*Blinov et al. in prep.*

→ all classes can “rotate” (HSP/LSP, FSRQs/BL Lacs, TeV and non-TeV)

▶ there is some dependence on the synchrotron peak with LSP rotations more often

→ both senses of rotation are allowed in the same source

▶ the rate can vary a lot for the same source



## EVPA rotations


*Blinov et al. 2015, MNRAS.453.1669B; Blinov et al. in prep.*

- ➔ all “rotators” are GL:
  - physical relation between  $\gamma$ -ray and optical polarization variability
- ➔ MC simulations: it is unlikely ( $p \leq 1.5 \times 10^{-2}$ ), that all the rotations are due to a random walk process

## EVPA rotations

*Blinov et al. 2015, MNRAS.453.1669B; Blinov et al. in prep.*

- data suggest:
  - ▶ the highest amplitude  $\gamma$ -ray flares are associated with smaller-than-average time lags
- two physical mechanisms:
  - ▶ one results higher amplitude flares and EVPA rotations
  - ▶ the other may be RW processes producing smaller amplitude flares, not related with rotations



## summary:

- high cadence, high precision optical linear polarization monitoring
- GL sources significantly more polarised:
  - ▶ *B*-field uniformity
  - ▶ non-thermal variability dominance
- 27 rotations found in 2 seasons (16 before RoboPol)
  - ▶ not all rotations are associated with a HE outburst
  - ▶ all “rotators” are GL: physical connection with  $\gamma$ -ray activity
  - ▶ unlikely that all are due to a random walk
  - ▶ data suggest: the highest amplitude  $\gamma$ -ray flares are associated with smaller-than-average time lags

# thank you

**Emmanouil Angelakis**

Max-Planck-Institut für Radioastronomie, Auf dem Huegel 69, Bonn 53121, Germany



MAX-PLANCK-GESELLSCHAFT



Max-Planck-Institut  
für Radioastronomie

