



Fermi

Gamma-ray Space Telescope

*Fermi* LARGE AREA  
TELESCOPE  
OBSERVATIONS OF  
HIGH-ENERGY  
GAMMA-RAY EMISSION  
FROM SOLAR FLARES

Melissa Pesce-Rollins,  
N. Omodei, V. Petrosian,  
W. Liu, F. R. da Costa,  
A. Allafort, L. Kashapova,  
V. Pal'shin

INFN-Pisa and Stanford HEPL  
[melissa.pesce.rollins@pi.infn.it](mailto:melissa.pesce.rollins@pi.infn.it)

on behalf of the *Fermi*-LAT  
collaboration

Fermi Symposium 2015


# THE *Fermi* SPACE TELESCOPE

## Gamma-ray Burst Monitor (GBM)


- ▶ 12 NaI and 2 BGO detectors
- ▶ Energy range: 8 keV–40 MeV
- ▶ Observes entire unocculted sky

## The Large Area Telescope (LAT)

- ▶ Pair conversion telescope
- ▶ Energy range: 20 MeV–> 300 GeV
- ▶ Large field of view ( $\approx 2.4$  sr): 20% of the sky at any time, all parts of the sky for 30 minutes every 3 hours
- ▶ Observes the Sun for  $\sim 20 - 40$  min every 3 hours




# WHY STUDY SOLAR FLARES WITH *Fermi*?



- ▶ Over the past 30 years limited sampling of solar flares with  $E > 25$  MeV
  - ▶ All of which were classified as GOES X class flares
  - ▶ Extended  $>100$  MeV emission for  $\sim 8$  hours detected by EGRET
  - ▶ 3 behind-the-limb flares with  $E < 100$  MeV


# WHY STUDY SOLAR FLARES WITH *Fermi*?



- ▶ Over the past 30 years limited sampling of solar flares with  $E > 25$  MeV
  - ▶ All of which were classified as GOES X class flares
  - ▶ Extended  $> 100$  MeV emission for  $\sim 8$  hours detected by EGRET
  - ▶ 3 behind-the-limb flares with  $E < 100$  MeV
- ▶ *Fermi* has detected more than 40 Solar flares in first 7 years of mission
  - ▶ More than half are GOES M class
  - ▶ Extended  $> 100$  MeV emission for more than 20 hours
  - ▶ Including 3 behind-the-limb flares with  $> 100$  MeV emission




# WHAT HAVE WE LEARNED SO FAR?



- ▶  $>100$  MeV lasting  $\sim$ hours from moderate GOES class flares is fairly common
- ▶ 13 hours of emission from the M3.7 GOES flare of 2011 March 7


*Fermi*  
Gamma-ray  
Space Telescope

# WHAT HAVE WE LEARNED SO FAR?



- ▶  $>100$  MeV lasting  $\sim$ hours from moderate GOES class flares is fairly common
- ▶ 13 hours of emission from the M3.7 GOES flare of 2011 March 7
  - ▶ Emission is well described by curved spectrum
  - ▶ Pion-decay is the most likely origin

# WHAT HAVE WE LEARNED SO FAR?




- ▶  $>100$  MeV lasting  $\sim$ hours from moderate GOES class flares is fairly common

- ▶ 13 hours of emission from the M3.7 GOES flare of 2011 March 7



- ▶ Emission is well described by curved spectrum

- ▶ Pion-decay is the most likely origin




- ▶  $>100$  MeV emission centroid consistent with active region for most bright flares

# WHAT HAVE WE LEARNED SO FAR?



Paper in preparation



- ▶  $>100$  MeV lasting  $\sim$ hours from moderate GOES class flares is fairly common

- ▶ 13 hours of emission from the M3.7 GOES flare of 2011 March 7


- ▶ Emission is well described by curved spectrum

- ▶ Pion-decay is the most likely origin

- ▶  $>100$  MeV emission centroid consistent with active region for most bright flares

- ▶ All LAT flares associated with fast CME's

# WHY ARE BEHIND-THE-LIMB FLARES INTERESTING?




Cliver et al. 1993


- ▶  $\gamma$ -ray emission processes require densities greater than  $10^{12} \text{ cm}^{-3}$
- ▶ Measurements of  $\gamma$ -ray line emission are generally consistent with a compact region located in the chromosphere
- ▶ Observations of  $\gamma$ -rays (both line emission and pion produced) from behind-the-limb flares can imply
  - ▶ A spatially extended flare component that can subtend a large range of heliolongitudes
    - ▶ Allowing the particles to interact at the visible disk
  - ▶ Or acceleration and emission occur in the Corona
    - ▶ Requires larger than usual Coronal densities

# LAT BEHIND-THE-LIMB FLARES


SOL2013-10-11



SOL2014-01-06




SOL2014-09-01




- ▶ Located  $\sim 10^\circ$  behind the eastern limb
- ▶  $>100$  MeV emission for 30 minutes
- ▶ *RHESSI* emission consistent with loop top
- ▶ Upper limits on nuclear line emission from GBM
- ▶ LAT emission centroid consistent with on-disk
- ▶ Located  $\sim 10^\circ$  behind the western limb
- ▶  $>100$  MeV emission for 20 minutes
- ▶ Associated with very strong SEP event
- ▶ Gamma-ray onset time consistent with Solar Particle Release time
- ▶ Insufficient statistics for localization
- ▶ Located  $\sim 40^\circ$  behind the eastern limb
- ▶  $>100$  MeV emission for  $\sim 2$  hours
- ▶ GBM and *Konus* emission up to few MeV
- ▶ 2.23 MeV line visible in GBM
- ▶ LAT emission centroid consistent with on-disk

# SPECTRAL ENERGY DISTRIBUTION OF SOL2013-10-11



- ▶ LAT spectra is curved – increase in significance of  $\sim 18\sigma$ 
  - ▶ Best proton spectral index is  $4.3 \pm 0.1$
- ▶ Only upper limits on the nuclear lines and neutron capture from GBM
- ▶ Broken power-law fit to  $<100$  keV GBM emission
  - ▶ Photon index  $3.22 \pm 0.005$
  - ▶  $E_{break} = 20 \pm 8$  keV

# SOL2014-09-01 SPECTRAL ENERGY DISTRIBUTION



1. Pion decay template
2. 2.23 MeV line (gaussian)
3. Power-law exponential cut-off
4. Power-law exponential cut-off

- ▶ LAT data is well described by a curved spectrum
  - ▶ Best proton spectral index from fit  $4.4 \pm 0.1$
- ▶ Power-law (3) index  $1.4 \pm 0.1$ , power-law (4) index  $2.7 \pm 0.1$
- ▶ Folding energy  $8.2 \pm 0.2$  MeV
- ▶ Significance of 2.23 MeV line is  $\sim 4\sigma$



# SUMMARY

- ▶ The *Fermi*-LAT has detected high energy gamma-rays from more than 40 solar flares over the first 7 years of mission
  - ▶ Almost half of which are GOES M class
  - ▶ Sampling both impulsive and sustained emission
  - ▶ Extended emission lasting hours is fairly common
  - ▶  $>100$  MeV emission is most likely due to pion-decay
- ▶ First detection of  $>100$  MeV emission from behind-the-limb Solar flares
  - ▶ Flares originate from behind both eastern and western limbs
  - ▶ Photons with energies up to 3 GeV measured from two of these flares
  - ▶ And 2.2 MeV neutron capture line from SOL2014-09-01
- ▶ Behind-the-limb flare observations seem to suggest a spatially extended component for high-energy gamma-rays
  - ▶ This component must subtend more  $30^\circ$  heliolongitude
  - ▶ Coronal Mass Ejection (CME) generated shocks could accelerate the particles over such a large range
  - ▶ Paper on LAT behind-the-limb flares in preparation

A large, light blue stylized logo of the Fermi Gamma-ray Space Telescope, featuring a curved tube and a central circular element with concentric rings.


SPARE SLIDES

*fermi*  
Gamma-ray  
Space Telescope

A total of 3 behind-the-limb flares were detected with emission below 100 MeV by space based observatories


1. 1989 September 29 flare detected by GRS on SMM
  - ▶ Vestrland & Forrest 1993, Cliver et al., 1993
  - ▶ Intense gamma-ray line emission in the 1-8 MeV range
  - ▶ Strong 2.2 MeV neutron capture line
  - ▶ Observations required spatially extended flare loop
2. 1991 June 1 flare detected by PHEBUS on GRANAT
  - ▶ Barat et al. 1994, Ramaty et al. 1997
  - ▶ Intense gamma-ray line emission in the 1-8 MeV range
  - ▶ No 2.2 MeV neutron capture line detected
  - ▶ Coronal origin for the emission was concluded
3. 1991 June 30 flare detected by PHEBUS, BATSE and EGRET
  - ▶ Vilmer et al. 1999, Trottet et al. 2003
  - ▶ No detectable line emission or 2.2 MeV line
  - ▶ Emission detected up to almost 100 MeV
  - ▶ Debate on Coronal or/and extended flare loop emission

# FIRST $>100$ MEV BEHIND THE LIMB FLARE



- ▶ Estimated GOES class from STEREO EUV emission is M4.9
- ▶ *RHESSI* and GBM detected emission up to 50 keV above the chromospheric limb
- ▶  $>100$  MeV emission detected for 25 minutes by LAT
- ▶ Pass7\_REP data published in ApJL, 805, L15
- ▶ Re-analyzed the flare with new *Fermi*-LAT Pass 8 data
  - ▶ Gained 5 minutes of detection with respect to Pass7\_REP
  - ▶ Detection from 07:10 - 07:40 UT

# TESTING THE EMISSION MODELS




We fit the LAT spectral data between 60 MeV and 10 GeV to test three different emission models:


1. Pure power-law
2. Power-law with exponential cut-off
3. Templates to describe emission from pion decay based on Murphy et al. 1987

We rely on the likelihood ratio test (TS) to estimate the significance of the source and whether the curved model provides a better fit



- ▶ When model (2) provides a better fit we also fit the data with a series of pion-decay models to determine the best proton spectral index



- ▶ On Jan 6, 2014 07:42 a flare erupted from an active region located S8W110
  - ▶ 20° behind the western limb
- ▶ A fast CME (speed  $\sim 1400$  km/s) was reported by LASCO
- ▶ Filament eruption detected from the visible limb by SDO
- ▶ RHESSI 6-12 keV source located above the limb
- ▶ LAT statistics insufficient to provide localization information




- ▶ Estimated GOES class based on STEREO EUV emission is X3.5
- ▶ Konus detected emission up to 78 keV
- ▶ *RHESSI* detected emission after 8:20 UT
- ▶ *Fermi* satellite was in the SAA from 7:25 - 7:55 UT
- ▶ Both detectors on-board *Fermi* detected emission from this flare upon exiting the SAA:
  - ▶ GBM detected emission in the 10's of keV range
  - ▶ LAT detected  $>100$  MeV emission for  $\sim 20$  minutes



- ▶ Very strong SEP event
- ▶ Use SEP onset times to estimate the acceleration time at the Sun
- ▶ Fit with a straight line assuming 1.2 AU propagation length gives 07:44 UT on Jan 6, 2014 ( $\pm 3$  minutes)
- ▶ In agreement with the Solar particle release time 07:47 UT reported by Thakur et al. ApJL 790, L13
- ▶ LAT detection starts 8 minutes later
  - ▶ *Fermi* in SAA until 7:55 UT
  - ▶ SEP and  $\gamma$ -producing ions accelerated at the same time?



# SOL2014-09-01 SPECTRAL ENERGY DISTRIBUTION




1. Pion decay template
2. 2.2 MeV line (gaussian)
3. Narrow lines template
4. Power-law with EXP cut-off


- ▶ LAT data is well described by a curved spectrum
  - ▶ Best proton spectral index from fit  $4.4 \pm 0.1$
- ▶ Photon spectral index  $2.1 \pm 0.1$
- ▶ Folding energy  $6.7 \pm 0.4$  MeV
- ▶ Significance of 2.23 MeV line and narrow lines is  $\sim 7\sigma$

# STEREO AND GOES PROTONS

SOL2013-10-11



SOL2014-09-01



Proton intensity-time profiles provide information on magnetic connectivity and SEP propagation



- ▶ SOL2013-10-11

- ▶ Active region located  $10^\circ$  behind eastern limb
- ▶ Poorly connected to Earth – no GOES signal following the flare

- ▶ SOL2014-09-01



- ▶ Active region located  $36^\circ$  behind eastern limb
- ▶ Protons reach Earth  $\sim 9$  hours after flare

# FINDING THE ONSET TIMES




- ▶ For GOES SEP we apply a median filtering algorithm to help smooth out the data
- ▶ Take the onset time to be the point of 5% max intensity
- ▶ We scan over a series of values for the median filter window
- ▶ Take median window 25 to be the onset time
- ▶ Take the difference in times over the scan values to be the error associated with the onset time

# FINDING THE ONSET TIMES FOR HE



- ▶ For GOES HE SEP we run an FFT on the data
- ▶ Take the onset time to be the point where second derivative is max
- ▶ We scan over a series of values for the frequency threshold
- ▶ Take frequency threshold of 13 to be the onset time
- ▶ Take the difference in times over the scan values to be the error associated with the onset time



## Toy model simulations

- ▶ Spherical harmonics expansion of the magnetic field ( $R < R_{\odot}$ )
  - ▶ First proposed by Newkirk et al. 1968
- ▶ Parker field for  $R > R_{\odot}$ 
  - ▶ Parker 1958, Giacalone & Jokipi 2004
- ▶ Particle propagation
  - ▶ Ray tracing of particles in magnetic field
  - ▶ Developing code to take into account the effects of scattering and turbulence

# FERMI LAT SOLAR FLARE PUBLIC DATA

Archive Search of Catalog(s)

Choose Tables > **Parameter Search** > Search Results > Choose Data Products

Description Catalog Data Default Radius (arcmin) Mission Table Type  
[fermi\\_lat\\_solar\\_flare\\_grb\\_catalog](#) Version: 1 ID: FERMI Object

1. Enter any constraints on the query below. [Help on constraint syntax](#)  
 (Help about [columns](#), [operators](#), and [sort ordering](#).)

2. To change the fields that are returned, select the box in the 'View' column beside each field desired.

3. To sort the results by any field, select one box in the 'Sort' column beside the field to sort on. [Examples of query constraints](#)

View	Sort	Parameter	Units	Query Syntax	Min Value	Max Value	Value Type
<input type="checkbox"/>	<input type="checkbox"/>	ISSAKI_ANGLE			0.0000000000	64.03027328	float
<input type="checkbox"/>	<input type="checkbox"/>	ISSAKI_SIZE			0.0000000000	31.6481233235	float
<input type="checkbox"/>	<input type="checkbox"/>	ISSAKI_TIME			00:47:12.0	23:33:36.2	position
<input type="checkbox"/>	<input type="checkbox"/>	ISSAKI_ANGLE			46:19:26	+75:51:23	position
<input type="checkbox"/>	<input type="checkbox"/>	ISSAKI_ANGLE			0.00000	337.8605	float
<input type="checkbox"/>	<input type="checkbox"/>	ISSAKI_ANGLE			65.1187	79.7746	float
<input type="checkbox"/>	<input type="checkbox"/>	ISSAKI_ANGLE			2008-08-25 13:57:08.185	2033-04-27 07:30:28.420	date
<input type="checkbox"/>	<input type="checkbox"/>	ISSAKI_ANGLE			2008-08-25 14:39:28.185	2033-04-27 08:03:41.420	date
<input type="checkbox"/>	<input type="checkbox"/>	ISSAKI_ANGLE			2008-08-25 14:13:48.105	2033-04-27 07:47:49.420	date
<input type="checkbox"/>	<input type="checkbox"/>	ISSAKI_ANGLE			GRB	GRBNAME	string
<input type="checkbox"/>	<input type="checkbox"/>	ISSAKI_ANGLE			0	0	integer

4. Do you want to change your current query settings?

Object Name Or Coordinates:  (e.g. Cyp X3 or 12 00 00, +12 0) Use semicolons(;) to separate multiple

Coordinates System:

Search Radius:  (arcmin) Default uses the optimum radius for each catalog searched.

Name Resolves:  (GRB, EPHEC, also NEI: g)

Observation Dates:  (Not all tables have observation dates. For those that do, the time portion of constraints with semicolons(;) Range separator: '..' (e.g. 1992-12-31, 48980.5; 2095-01-31, 12 00:00.39)

Limit Results To:  rows

Output Format:


Show All Parameters:  Select to display all catalog parameters instead of only defaults

5.

## Fermi LLE public data

- ▶ LLE catalog of Solar flares and GRBs
- ▶ 11 impulsive solar flares and 56 GRBs
- ▶ <http://heasarc.gsfc.nasa.gov/W3Browse/fermi/fermille.html>
- ▶ All LLE data products publicly available
  - ▶ LLE event file
  - ▶ spectrum files (PHAI, PHAI and RSP)
  - ▶ Quick look files
- ▶ LLE data can be analyzed with XSPEC and rmfit

March 7, 2011 M3.7 class flare



<http://sprg.ssl.berkeley.edu/~tohan/brower>

## Fermi LAT SunMonitor

- ▶ Fermi-LAT SunMonitor continuously monitors the Sun
- ▶ >100 MeV gamma-ray flux from the Sun in fixed 3 hour time intervals
- ▶ All available online!