Lower and Upper Bounds on the Cosmic TeV Gamma-ray Background

Yoshiyuki Inoue

(ISAS/JAXA, JAXA International Top Young Fellow) Collaborators: Yasuyuki T. Tanaka



6th Fermi Symposium, Washington D.C., 2015-11-11

Cosmic Gamma-ray Background Spectrum at >0.1 GeV



 Fermi has resolved 30% of the CGB at ~1 GeV and more at higher energies (see Di Mauro's talk).



Components of the Cosmic GeV Gamma-ray Background



- Blazars (Ajello+'15), Radio gals. (YI'11), & Star-forming galaxies (Ackermann+'12) make up almost 100% of CGB from 0.1-1000 GeV.
- Next frontiers will be
 - Anisotropy (e.g. Ando & Komatsu '06, Ackermann+'11, Camero+'13, Shirasaki+'14)
 - Cosmic MeV Gamma-ray Background (e.g. YI+'08, Ajello+'09, YI+'13)
 - <u>Cosmic TeV Gamma-ray Background (This talk)</u>

Cosmic TeV Gamma-ray Background



- Above 1 TeV, there is no gamma-ray data, though it is important for neutrino studies.
- extragalactic pp scenario for IceCube events is constrained by the CGB (Murase+'13; Bechtol+'15).

Cosmic TeV Gamma-ray Background



 Above 1 TeV, there is no gamma-ray data, though it is important for neutrino studies.

 extragalactic pp scenario for IceCube events is constrained by the CGB (Murase+'13; Bechtol+'15).

GeV-TeV Gamma-ray Connection: Cascade



- TeV gamma-ray photons are absorbed by EBL
- electron-positron pairs are created
- pairs scatter CMB via inverse-Compton process
 - 1 TeV (primary) -> ~1 GeV (secondary)
- Note: plasma instability may suppress the cascade (Broderick+'12, but see also Sironi & Giannios '14)

Upper Bound on the Cosmic Gamma-ray Background



- Cascade component from the TeV background can not exceed the Fermi data (Coppi & Aharonian '97, YI & Ioka '12, Murase+'12, Ackermann+'14).
 - No or negative evolution is required -> low-luminosity BL Lacs show negative evolution (Ajello+'14).

Galaxy Counts: Lower Bound on the Cosmic Optical/Infrared Background

TeV blazar sample

 Select 36 blazars the default TeVcat catalog.

- Low-state data are available for 31/36.
- 3FGL SED data for the GeV data.
- Radio galaxies and star-forming galaxies are not included yet.

Lower Bound on the Cosmic Gamma-ray Background

TeV blazar counts give lower limit on to the cosmic gamma-ray background.

- Fermi has resolved more portion of the TeV sky than IACTs do?
 - CTA survey will be important (YI, Totani, & Mori 10; Dubus, YI, +'13)

Bounds on the Cosmic TeV Gamma-ray Background

Current limit at 0.3-10 TeV is

$$3 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-1} \left[\text{MeV/cm}^2/\text{s/sr}\right] < E^2 \frac{dN}{dE} < 5 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-0.7} \left[\text{MeV/cm}^2/\text{s/sr}\right]$$

Bounds on the Cosmic TeV Gamma-ray Background

Current limit at 0.3-10 TeV is

$$3 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-1} \left[\text{MeV/cm}^2/\text{s/sr}\right] < E^2 \frac{dN}{dE} < 5 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-0.7} \left[\text{MeV/cm}^2/\text{s/sr}\right]$$

Bounds on the Cosmic TeV Gamma-ray Background

Current limit at 0.3-10 TeV is

$$3 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-1} \left[\text{MeV/cm}^2/\text{s/sr}\right] < E^2 \frac{dN}{dE} < 5 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-0.7} \left[\text{MeV/cm}^2/\text{s/sr}\right]$$

- Cosmic TeV gamma-ray background is not well investigated yet.
- Current GeV gamma-ray background gives upper limits on the TeV gamma-ray background through the cascade argument
- Ensemble of low-state TeV blazar flux gives lower limit on to the cosmic gamma-ray background
- Current limit on the TeV background is

 $3 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-1} \, \left[\text{MeV/cm}^2/\text{s/sr}\right] < E^2 \frac{dN}{dE} < 5 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-0.7} \, \left[\text{MeV/cm}^2/\text{s/sr}\right]$

Lower Bound on the Cosmic Gamma-ray Background

TeV blazar counts give lower limit on to the cosmic gamma-ray background.

Current limit at 0.3-10 TeV is

$$3 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-1} \, \left[\text{MeV/cm}^2/\text{s/sr}\right] < E^2 \frac{dN}{dE} < 5 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-0.7} \, \left[\text{MeV/cm}^2/\text{s/sr}\right]$$

Fermi has resolved more portion of the TeV sky than IACTs do?

CTA survey will be important (YI, Totani, & Mori 10; Dubus, YI, +'13)

Lower Bound on the Cosmic Gamma-ray Background

TeV blazar counts give lower limit on to the cosmic gamma-ray background.

Current limit at 0.3-10 TeV is

$$3 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-1} \left[\text{MeV/cm}^2/\text{s/sr}\right] < E^2 \frac{dN}{dE} < 5 \times 10^{-5} \left(\frac{E}{100 \text{GeV}}\right)^{-0.7} \left[\text{MeV/cm}^2/\text{s/sr}\right]$$

Fermi has resolved more portion of the TeV sky than IACTs do?

CTA survey will be important (YI, Totani, & Mori 10; Dubus, YI, +'13)

VHE EGB Model

$$E^{2} \frac{dN}{dE} (E_{\text{obs}}) = \frac{cE_{\text{obs}}^{2}}{4\pi} \int_{0}^{z_{\text{max}}} dz \left| \frac{dt}{dz} \right| (1+z)$$
$$\times \frac{dj}{dE_{\gamma}} [(1+z)E_{\text{obs}}, z] \exp[-\tau_{\gamma\gamma}(E_{\text{obs}}, z)],$$
$$j = j_{\text{int}} + j_{\text{cas}} \qquad \text{MeV/s/MeV/cm}^{3}$$

$$\frac{dj_{\text{int}}}{dE_{\gamma}}(E_{\gamma},z) = \begin{cases} j_0 E_{\gamma}^{-\Gamma_{\text{ph}}} (1+z)^{\beta_{\text{evo}}}, & E_{\gamma} \leq E_{\text{max}}, \\ 0, & E_{\gamma} > E_{\text{max}}, \end{cases}$$

- Parameters are
 - β_{evo} : Cosmological evolution of number density
 - Most of known sources show $\beta_{evo} > 0$
 - E_{max} : Maximum energy
 - Γ_{ph}: Photon index

Upper Limit on EGB

- $\beta_{evo}=0$ (no evolution)
- Γ_{ph}=1.5 (Fermi acc. limit)
- $E_{max} = 60 \text{ TeV}$
- Normalization is fixed to the EGB data < 100 GeV.
- No known source classes included.
- Upper limit is consistent with the Fermi VHE EGB obs.

• UL:
$$E^2 \frac{dN}{dE} < 1.1 \times 10^{-4} \left(\frac{E}{100 \,\text{GeV}}\right)^{-0.5} \text{MeV/cm}^2/\text{s/sr}$$

Different Photon Indices

- Γ~1.5 is the most conservative.
- Harder spectral model is limited by the cascade.
- Softer spectral model is limited by the primary.

Different E_{max} and evolution

- β >0 violates the limit.
- β <0 eases the limit.
 - But, no known gammaray sources.
- low E_{max}(~sub-TeV) eases the limit.
- If cosmological,
 - sources should have a hard spectrum and show no or negative evolution.

Components of EGB

 FSRQs (Ajello+'12), BL Lacs (Abdo+'10), Radio gals. (YI'11), Starburst gals. (Stecker & Venters '11, Ackermann+'12) are guaranteed to contribute to EGB.

in mindWe need to subtract them to evaluate the VHE EGB upper lighterror band

Upper Limit on EGB w/ known sources

If we try to explain EGB at <10GeV, the observation violates the limit.

• UL:
$$E^2 \frac{dN}{dE} < 4.5 \times 10^{-5} \left(\frac{E}{100 \,\text{GeV}}\right)^{-0.7} \text{MeV/cm}^2/\text{s/sr}$$

Upper Limit on EGB w/ other models

If we try not to violate the limit, residual appears at <10GeV.

Possible Explanations

- Hard spectrum with sub-TeV E_{max} and $\beta{<}0$
 - No known sources. TeV HBL? Low-luminosity GRBs?
- New Physics: Axion or Lorentz invariance violation?
- Dark matter in local?
- ~GeV sources
 - pulsars? radio-quiet AGNs?
- Foreground uncertainty?
- See also Murase+'12

