



# Active Galactic Nuclei with Fermi-LAT

Elisabetta Cavazzuti  
ASI and GSFC  
(with the help of past LAT speakers)  
Fermi Summer School 2017, Lewes



# What is an Active Galactic Nuclei

# Active Galactic Nuclei

---

- ★ Objects, sometimes **looking like galaxies**, other times **apparently stellar**, which show **extreme amount of radiation**, and sometimes powerful jets of material from deep in their centers

# Active Galactic Nuclei

---

- ★ Objects, sometimes **looking like galaxies**, other times **apparently stellar**, which show **extreme amount of radiation**, and sometimes powerful jets of material from deep in their centers
- ★ Brightness can change significantly in several months/days/hours so the size must be very small

# Active Galactic Nuclei

---

- ★ Objects, sometimes **looking like galaxies**, other times **apparently stellar**, which show **extreme amount of radiation**, and sometimes powerful jets of material from deep in their centers
- ★ Brightness can change significantly in several months/days/hours so the size must be very small
- ★ All the various AGN types: accretion of matter onto the central super-massive black hole (~billion solar mass)
  - ★ When there is accretion: we have an AGN
  - ★ When there is not accretion: AGN is dormant and galaxy looks normal

# Active Galactic Nuclei

---

- ★ Objects, sometimes **looking like galaxies**, other times **apparently stellar**, which show **extreme amount of radiation**, and sometimes powerful jets of material from deep in their centers
- ★ Brightness can change significantly in several months/days/hours so the size must be very small
- ★ All the various AGN types: accretion of matter onto the central super-massive black hole (~billion solar mass)
  - ★ When there is accretion: we have an AGN
  - ★ When there is not accretion: AGN is dormant and galaxy looks normal
- ★ All AGN types are a combination of two phenomena:
  - ★ Amount of accretion onto the SMBH: **LUMINOSITY**
  - ★ Orientation angle of the galaxy/AGN respect to the observer: **AGN type**

# Active Galactic Nuclei


---

- ★ Objects, sometimes **looking like galaxies**, other times **apparently stellar**, which show **extreme amount of radiation**, and sometimes powerful jets of material from deep in their centers
- ★ Brightness can change significantly in several months/days/hours so the size must be very small
- ★ All the various AGN types: accretion of matter onto the central super-massive black hole (~billion solar mass)
  - ★ When there is accretion: we have an AGN
  - ★ When there is not accretion: AGN is dormant and galaxy looks normal
- ★ All AGN types are a combination of two phenomena:
  - ★ Amount of accretion onto the SMBH: **LUMINOSITY**
  - ★ Orientation angle of the galaxy/AGN respect to the observer: **AGN type**

**Up to many thousand times more luminous than the entire Milky Way but the energy is released within a region approximately the size of our solar system!**

# Active Galactic Nuclei


Urry&Padovani 1995






# Active Galactic Nuclei

Urry&Padovani 1995




# Active Galactic Nuclei

Urry&Padovani 1995




# Active Galactic Nuclei

Urry&Padovani 1995




# Active Galactic Nuclei

Urry&Padovani 1995




# Active Galactic Nuclei

Urry&Padovani 1995




# Active Galactic Nuclei

Urry&Padovani 1995



# Active Galactic Nuclei

Urry&Padovani 1995




## Radio loudness:

$$\frac{\text{Radio (5GHz) Flux}}{\text{Optical (B) Flux}} \geq 10$$

(Kellermann et al. 1989)

# Active Galactic Nuclei

Urry&Padovani 1995



**Radio-loud objects** have emission contributions from both the jet(s) and the lobes that the jets inflate.

These emission contributions dominate the luminosity of the AGN at nearly all  $\lambda$


### Radio loudness:

$$\frac{\text{Radio (5GHz) Flux}}{\text{Optical (B) Flux}} \geq 10$$

(Kellermann et al. 1989)

**Radio-quiet objects** are simpler since jet and any jet-related emission can be neglected at all wavelengths.





**Radio-quiet objects** are simpler since jet and any jet-related emission can be neglected at all wavelengths.

**AGN are a few % of all galaxies**

**Radio-loud objects** have emission contributions from both the jet(s) and the lobes that the jets inflate. These emission contributions dominate the luminosity of the AGN at nearly all  $\lambda$

### Radio loudness:

$$\frac{\text{Radio (5GHz) Flux}}{\text{Optical (B) Flux}} \geq 10$$

(Kellermann et al. 1989)

- Central nucleus outshines the rest of the galaxy
- High luminosity
- Emission across the entire e.m. spectrum, from radio to TeV
  - > non thermal
- Strong variability
- Radio-loud sources
  - > relativistic jets, superluminal motion

# Radio Galaxies






## 5-15% of active galaxies

(others are mostly Seyferts)


- ✓ Mostly of the elliptical type (Seyferts : spirals)
- ✓ Galaxies displaying extended radio lobes (up to 10 x larger than the galaxy)

### 2 classes

- **Fanaroff-Riley 1:** large opening angle, brighter close to the core, low luminosity, close
- **Fanaroff-Riley 2:** highly collimated jet, lobe brightened with hot spots, luminous, distant




# Blazars



< 5% of the whole AGN class

- ✓ beamed (jet at  $\leq 20\text{-}30^\circ$ ) (point at us!)
- ✓ broad band SED,  
L up to  $10^{49}$  erg  $\text{s}^{-1}$
- ✓ compact morphology  
(core flux  $\gg$  extended flux)
- ✓ flat radio spectrum ( $\alpha_r \leq 0.5$ )
- ✓ rapid and large variability (large  $\Delta L / \Delta t$ ),  
superluminal motions
- ✓ high and variable optical polarization

# Blazars




< 5% of the whole AGN class

- ✓ beamed (jet at  $\leq 20\text{-}30^\circ$ ) (point at us!)
- ✓ broad band SED,
  - L up to  $10^{49}$  erg  $\text{s}^{-1}$
- ✓ compact morphology
  - (core flux  $\gg$  extended flux)
- ✓ flat radio spectrum ( $\alpha_r \leq 0.5$ )
- ✓ rapid and large variability (large  $\Delta L / \Delta t$ ),
  - superluminal motions
- ✓ high and variable optical polarization

blazars dominate the extragalactic sky in a number of *new* observational windows  
( $\mu$ -wave, hard X-ray,  $\gamma$ -ray, TeV)

# Blazars




< 5% of the whole AGN class

- ✓ beamed (jet at  $\leq 20\text{-}30^\circ$ ) (point at us!)
- ✓ broad band SED,  
L up to  $10^{49}$  erg  $\text{s}^{-1}$
- ✓ compact morphology  
(core flux  $\gg$  extended flux)
- ✓ flat radio spectrum ( $\alpha_r \leq 0.5$ )
- ✓ rapid and large variability (large  $\Delta L / \Delta t$ ),  
superluminal motions
- ✓ high and variable optical polarization


blazars dominate the extragalactic sky in a number of *new* observational windows  
( $\mu$ -wave, hard X-ray,  $\gamma$ -ray, TeV)

As of today, about **3.561 blazars** (**BZCAT 5<sup>th</sup> edition**, [www.asdc.asi.it/bzcat](http://www.asdc.asi.it/bzcat)) are known and before the operations of Fermi, Swift, Planck and other facilities, only small good statistical samples existed.

## What do they look like?




## What do they look like?



Credit: SDSS /  
Size: 3' × 3'

# What do they look like?





## Blazar classes

**FSRQ:** Flat Spectrum Radio Quasar

**BL Lac:** named after prototype BL Lacertae

class	FSRQs	BLLacs
Defining property	Optical spectrum: strong broad emission lines	Optical spectrum: nearly featureless
Environment	intense radiation field (disk, clouds, torus)	low radiation field
Power	$\sim 10^{46-48}$ erg/s	$\sim 10^{45-46}$ erg/s
Parent population	Fanaroff-Riley 2	Fanaroff-Riley 1
Synchrotron hump in SED	Peak in IR	Peak in Opt/IR: LBL Peak in UV/X-rays: HBL
Redshift	0.1 – 4...	< 1 ??????

## Blazar classes

**FSRQ:** Flat Spectrum Radio Quasar

**BL Lac:** named after prototype BL Lacertae

class	FSRQs	BLLacs
Defining property	Optical spectrum: strong broad emission lines	Optical spectrum: nearly featureless
Environment	intense radiation field (disk, clouds, torus)	low radiation field
Power	$\sim 10^{46-48}$ erg/s	$\sim 10^{45-46}$ erg/s
Parent population	Fanaroff-Riley 2	Fanaroff-Riley 1
Synchrotron hump in SED	Peak in IR	Peak in Opt/IR: LBL Peak in UV/X-rays: HBL
Redshift	0.1 – 4...	<b>&lt; 1 ??????</b>



**BL Lacs redshifts are hard to measure, even at Kech Telescope!  
Many redshifts are missing. Be careful when you draw conclusions!**

# Blazar classes

**FSRQ:** Flat Spectrum Radio Quasar

**BL Lac:** named after prototype BL Lacertae

class	FSRQs	BLLacs
Defining property	Optical spectrum: strong broad	Optical spectrum: nearly




# Blazar classes


**FSRQ:** Flat Spectrum Radio Quasar

**BL Lac:** named after prototype BL Lacertae


class	FSRQs	BLLacs
Defining property	Optical spectrum: strong broad	Optical spectrum: nearly



# Blazars: SED properties




# Blazars: SED properties



# Blazars: SED properties

“two-hump” SEDs in  $\nu F_\nu$


sed-1229p0203 Ra=187.27750 deg Dec=2.05228 deg (NH=1.7E20 cm<sup>-2</sup>)



# Blazars: SED properties

“two-hump” SEDs in  $\nu F_\nu$

sed-1229p0203 Ra=187.27750 deg Dec=2.05228 deg (NH=1.7E20 cm<sup>-2</sup>)





# Blazars: SED properties

Low synchrotron peaked

**LSP/LBL**

IR-optical ( $\nu_{\text{syn}} < 10^{14} \text{ Hz}$ )

High synchrotron peaked



**HSP/HBL**

X-rays ( $\nu_{\text{syn}} > 10^{15} \text{ Hz}$ )

Intermediate synchrotron peaked

**ISP/IBL**


UV ( $10^{14} < \nu_{\text{syn}} < 10^{15} \text{ Hz}$ )



# Fermi LAT AGN

# Third catalog of AGN with LAT

FSRQ, BL Lac, BCU




**The 3rd Catalog of AGN Detected by the Fermi LAT (3LAC)**

Ackermann, M. et al. 2015, ApJ, 810, 14

# AGN classification in LAC catalogs

Two classification schemes:

**Optically-based** (strength of broad lines): FSRQs, BL Lacs, BCUs (Blazar Candidate of Unknown type)





# AGN classification in LAC catalogs


Two classification schemes:

**Optically-based** (strength of broad lines): FSRQs, BL Lacs, BCUs (Blazar Candidate of Unknown type)

**SED-based**: Low-, Intermediate-, High-Synchrotron-Peaked sources (LSPs, ISPs, HSPs resp.)




## Flux and energy flux




The solid (dashed) curve represents the approximate 3FGL (2FGL) detection limit based on a typical exposure.

# Flux and energy flux




The solid (dashed) curve represents the approximate 3FGL (2FGL) detection limit based on a typical exposure.

**Strong bias in photon flux but not in energy flux and thus in luminosity neither**



# Luminosity





## Blazar population

---


**CAVEAT !!!**

**295/604 BL Lacs have no measured redshifts  
(55%, 61%, 40%) for (LSPs, ISPs and HSPs)**

**Form your own opinion about it!**


Blazar Sequence, Blazar Divide (Fossati, Ghisellini et al xxx)  
Giommi&Padovani (xxx)

## 3LAC and the missing FSRQ




Missing sources

- FGL catalogs detect less LSP sources than HSP ones.
- Steep sources (such as FSRQ) show the brightest gamma-ray flares.
- Such big flares allow to detect the steeper sources, otherwise not enough bright (in gamma-ray).




# Transient gamma-ray sources




Flat Spectrum – Low Synchrotron peaked radio source

# Transient gamma-ray sources



Flat Spectrum – Low Synchrotron peaked radio source

## A variable sky



CRATES source:  
Typical Flat Spectrum  
Low Synchrotron  
Peaked radio source

TS=91 in month 46  
TS=106 in month 55

**A certain Region of  
Interest, analysed over  
77 months**

- 3FGL
- 2FGL
- 1FGL
- CRATES
- BZCAT


## **Some (past and) recent major results**

# Radio galaxies


Local **radio galaxies** ( $z < 0.1$ ) and blazars occupy different regions of the plot, with non-blazar AGN generally characterized by lower luminosity.

On the contrary, the two more distant **steep spectrum radio quasars** ( $z > 0.6$ ) fall within the range of  $\gamma$ -ray luminosities of FSRQs.

The non-blazar AGN detected by Fermi so far are characterized by **large core dominances** (ratio between IC and Synch peak fluxes)




# Fornax A lobes




A leptonic emission that arises due to IC scattering of EBL photons off of relativistic electrons in the radio lobes underestimates the observed  $\gamma$ -ray emission for any current EBL estimate

A hadronic-only model (proton-proton interactions) requires implausibly large total cosmic-ray energy when compared to an estimate of the Fornax A outburst assumed to have created the lobes






## The most distant $\gamma$ -ray detected blazars



## The most distant $\gamma$ -ray detected blazars



NVSS J151002+570243 ( $z = 4.31$ ) is now the farthest known  $\gamma$ -ray emitting blazar


Ackermann, M. et al. 2017, ApJL, 837, L5

~1.4million quasars included in the Million Quasar Catalog (MQC; Flesch 2015)

- select all  $z > 3.1$  sources and
- retain only radio-loud
- 92 months of LAT data (ROI) of  $15^\circ$  radius centered on each quasar and define a sky model that includes all  $\gamma$ -ray sources detected in the 3FGL


expand the energy range considered so it spans **60 MeV to 300 GeV to be more sensitive to spectrally soft  $\gamma$ -ray sources**, i.e., blazars whose high-energy peak is shifted to lower energies ( $\sim 1$ – $10$  MeV) as typical for high- $z$  blazars.

# CTA 102 (FSRQ)




the  $\gamma$ -ray outburst is coincident with flares at all the other frequencies and is related to the passage of a new superluminal knot through the radio core.

Casadio et al, 2015ApJ, 813, 51C




# CTA 102 (FSRQ)



Casadio et al, 2015ApJ, 813, 51C

erratic blazar nature, revealing **both strong connections across wave bands in one outburst and no obvious connections for other events.**


# Quasi-periodic Modulation in PG 1553+113



- BL Lac HSP  
 $0.395 < z < 0.62$
- one of the brightest BL Lacs in the X-ray band
- TeV source (HESS, MAGIC)
- PeV  $\nu$  ?


$\sim 6.5$  years  
Pass8 data

# Quasi-periodic Modulation in PG 1553+113



Critical MW data

## Quasi-periodic Modulation in PG 1553+113



The periodicity analysis methods applied to optical and radio light curves found compliant results with the gamma-ray light curves.

- Composite optical light curve period search:  $\sim 752 \pm 8$  days ( $2.06 \pm 0.02$  years)
- Radio light curve period search:  $\sim 695 \pm 75$  days ( $1.9 \pm 0.2$  years)

## Contribution to Extragalactic Gamma-Ray Background


EGB total intensity of  $1.1 \times 10^{-5} \text{ ph cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$

Blazars contribute a grand-total of  $(5-7) \times 10^{-6} \text{ ph cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$

– Resolved sources :  $\sim 4 \times 10^{-6} \text{ ph cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$

– Unresolved blazars:  $\sim (2-3) \times 10^{-6} \text{ ph cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$  (in agreement with Abdo+10)

Blazars, star-forming galaxies and radio galaxies can explain the intensity and the spectrum of the EGB







## **GeV – TeV connection**

# Fermi High energy source Lists (FHL)


**1FHL > 10 GeV,  
36 months years**



**2FHL, > 50 GeV,  
80 months**




**3FHL, > 10 GeV  
84 months**



**AGN are still the vast majority of counterparts**

# Blazars synchrotron peak distributions



Different population than 3FGL

## Blazars synchrotron peak distributions




Fig. 14.— Normalized distributions of the frequency of the synchrotron peak for the blazars detected in the 3FGL (0.1–300 GeV), 2FHL (50 GeV–2 TeV), and 3FHL (10 GeV–2 TeV) catalogs.

## BL Lac spectral index distributions










Fig. 10.— The distribution of spectral indices for a subsample of 158 BL Lacs that are in common among the 2FHL (backslash orange), 3FGL (slash green), and 1FHL (purple). The medians of the distributions are shown with vertical lines. The higher the energy band, the larger the index; therefore sources get softer with increasing energy. The scatter of the distribution is also larger with increasing energy, partly because of the lower statistics.

# Multi wavelength high energy SED





# Multi wavelength high energy SED




Being sensitive over  $\sim 4$  decades in energy, the LAT resolves the high-energy peak

- Sources become softer at higher energies
- Sources becomes softer at high redshift



## EBL attenuation


Photons with energies greater than about 30 GeV suffer from attenuation over cosmological distances as a consequence of the pair production interactions with extragalactic background light (EBL) photons



clear softening of the spectral index above 10 GeV with increasing redshift, which is likely due to EBL attenuation.



# Photon spectral index



The trend of **a strong hardening of the energy spectra with increasing peak frequency** already seen above 100 MeV in the Fermi-LAT AGN catalogs (e.g., Ackermann et al. 2015) is even more pronounced above 10 GeV. This enhanced effect relative to 3LAC is due to the larger EBL attenuation suffered by high-redshift sources (most of them being LSPs) in comparison with the lower-redshift ones (preferentially HSPs).

**Thank you for listening!**