

2016 data: profile plots with best  
aligned geometry (so far)

Alessandra Filippi

May 14, 2018

# 2016 data @ 0.5 mm


## Moller and FEE analysis with Sho's cuts

- One good detector chosen (my reference: v5-21)
- In reconstruction: fixed beamspot (0,0) and  $z_{\text{target}}=0$ 
  - This means:  $z_{\text{vtx}}$  still wrong
- Purpose
  - analyse Moller pairs and FEE tracks with the same cuts applied by Sho for 2015 data (changing scale where needed)
  - Check consistency with 2015 results and internal consistency
- Input: FEE and Moller ntuple out of the reconstruction
  - Checked by Miriam: no double corrections
  - Outputs in mm! (while hps-java gets offsets in cm... a little misunderstanding of scale)


# Moller events: common cuts

- Top-bottom track time:
  - $|\text{topTrkT} - \text{botTrkT}| < 3 \text{ ns}$
- Tracks in detector acceptance (close to z axis):
  - Large  $\theta_x$  angle for tracks (from unconstrained and fitted momentum):
    - $|\text{uncPX}/\text{uncP}| < 0.005$
    - $|\text{topPX}/\text{topP}| < 0.01$  (same for bottom)
  - Large  $\theta_y$  angle (from unconstrained and fitted momentum):
    - $|\text{uncPY}/\text{uncP}| < 0.005$

# Moller evts: $z_0$ impact parameter vs $\cos\theta_y$




The TOP distribution has a less uniform structure (striped? Why only for top??)  
 Some troubles with y coordinate reconstruction? (could be due to strip pitch by why not seen in both halves?)






	Top	Bottom
P0	-0.12	0.32
P1	2.30	9.09

# Moller evts: $d_0$ impact parameter vs $\cos\theta_y$



No structures seen (the problem is in the  $y$  coordinate!)




	Top	Bottom
P0	-0.28	-0.52
P1	0.25	0.56

# Moller evts: $d_0$ impact parameter vs $\cos\theta_y$ in energy steps - TOP half

Selection in energy intervals 160 MeV wide from  $\sim 700$  MeV


A dependence on energy should not be desirable (this would imply a dependence on acceptance)... but there is

Some sort of parabolic trend of  $d_0$  central value




# Moller evts: $d_0$ impact parameter vs $\cos\theta_y$ in energy steps- BOTTOM half

Selection in energy intervals 160 MeV wide from  $\sim 700$  MeV




# Moller evts: invariant mass ( $e^-e^-$ ) vs $\Delta\phi$ (opening angle between the two tracks)



When the tracks are at large azimuthal angles, the invariant mass of the electron pair is larger



# Moller events: invariant mass ( $e^-e^-$ ) vs $\Delta p(\text{top-bottom})$




Flat enough to be happy enough

# $\theta_z - \theta$ (from Moller formula) vs $\varphi$ in energy ranges – TOP half

Trend of dip angle correction as a function of the azimuth angle and energy

$$y = \arccos \theta_z - \arccos(1 - m_e(1/p - 1/E_b))$$

$$x = \arctan(p_y, p_x)$$



Selection in energy intervals 160 MeV wide from  $\sim 700$  MeV



Flat enough



# FEE events: common cuts

- Trigger:
  - isSingle0 || isSingle1
- Max number of hits per track:
  - fspTrkHits==6
- Ecal-svt match  $\chi^2$ :
  - fspMatchChisq<3
- Ecal cluster energy: 85% Ebeam
  - fspCIE < 0.85\*Ebeam
- No cut of track fit quality (track  $\chi^2$  – I usually ask  $\chi^2 < 20$ )


# FEE: $z_0$ impact parameter vs $\cos\theta_y$



	Top	Bottom
P0	-0.17	0.17
P1	5.20	5.95


Use these values as  $z_{\text{Tar}}$  input  
for alignment?  
(the “old famous” 5 mm...)

# FEE: $d_0$ impact parameter vs $\cos\theta_x$



top


bottom



	Top	Bottom
P0	0.1	0.01
P1	5.98	4.82


# FEE: $p$ vs $\cos\theta_x$

Elastic electrons, top




top

Fitted value of par[1]=Mean




bottom

Elastic electrons, bottom




Fitted value of par[1]=Mean



	Top	Bottom
P0	2.26	2.26
P1	-0.63	-1.54


# FEE: $p$ vs $\cos\theta_y$

$p$




top


Electron side:  $\text{fspPX}/\text{fspP} < 0.01$   
(hole side)



$p$



bottom



	Top	Bottom
P0	2.39	2.93
P1	-3.76	-0.20



# ToDo list

- FEE plots are the cleanest and reliable ones
  - Results are consistent with what was found for 2015 data
  - Both  $z_0$  and  $d_0$  scatter plots indicate the  $z_{\text{Tar}}$  is at about -5 mm
- Try to use the information from these plots to fix the position of the target (use the values provided by the scatter plots as offsets and check results)
  - Inserting the  $z_{\text{tar}}$  information as millepede global offset for z translations of all sensors is not particularly useful, as this offset is absorbed by other z alignment corrections
  - Check the effect on the reconstruction if the new target position is inserted in ReconParticleDriver