

HPS-Sim Update

Bradley Yale


09/11/2018

Some hps-sim notes


- Hps-sim is much faster now, thanks to Maurik's optimization
 - Only a few hours on the farm, instead of days
- Hps-sim currently keeps far more particles than SLIC
 - Each hps-sim file is 2.4 GB! (vs. ~900 MB for SLIC)
- Things looked screwy in events mixed by hps-sim, so I checked stdhep-mixed events that were simply propagated with hps-sim.
- These comparisons had to be normalized by `n_events` instead of `n_files`

Wab-beam-tri events mixed using hps-sim (recon follow-up from last time)

Electron Momentum




Positron Momentum




Hps-sim mixed (Px)

Unconstrained Px




Target-Constrained Px




Hps-sim mixed (Py)

Unconstrained Py




Target-Constrained Py




Hps-sim mixed (Pz)

Unconstrained Pz




Target-Constrained Pz




Hps-sim mixed (x-vtx)

X-vtx (unconstrained)




X-vtx (target-constrained)




Hps-sim mixed (γ -vtx)

Y-vtx (unconstrained)




Y-vtx (target-constrained)




Hps-sim mixed (z-vtx)

Z-vtx (unconstrained)




Z-vtx (target-constrained)




Hps-sim mixed (mass)

Unconstrained Mass




Target-Constrained Mass




Using only stdhep-mixed events (same input)

Electron Momentum




Positron Momentum




Stdhep-mixed events (Px)

Unconstrained Px




Target-Constrained Px




Stdhep-mixed events (Py)

Unconstrained Py




Target-Constrained Py




Stdhep-mixed events (Pz)

Unconstrained Pz




Target-Constrained Pz




Stdhep-mixed events (mass)

Unconstrained Mass




Target-Constrained Mass




Stdhep-mixed events (x-vtx)

X-vtx (unconstrained)




X-vtx (target-constrained)




Stdhep-mixed events (γ -vtx)

Y-vtx (unconstrained)




Y-vtx (target-constrained)




Stdhep-mixed events (z-vtx)

Z-vtx (unconstrained)




Z-vtx (target-constrained)




2D Vertex locations (unconstrained)

uncVY:uncVX




uncVY:uncVX




2D Vertex locations (target-constrained)

tarVY:tarVX




tarVY:tarVX




Timing

Cluster Time diff.




Track Time diff.




Other plots

Momentum Asymmetry




Positron D0




Looking at DSTs (tri tuples previously shown)

Xvtx




Yvtx




Slic vs. Hps-sim (DST V0's)

ZVtx




Mass




Track-Cluster Matching

HitX-TrackX





HitY-TrackY



Momentum (DST V0's)

Px

Py




Summary


- Hps-sim problems seem mostly independent of its mixing procedure
- Hps-sim has missing WBT events at the target
 - Need to look at tritrig (simpler events) next
- More digging is needed

Momentum (DST V0's)

Pz



Track Slope



phi1-phi2

